

**Management in Practice****The first eradication of an exotic ant species from the entirety of Australia: *Pheidole fervens***Benjamin D. Hoffmann<sup>1</sup>, Jim Eldridge<sup>2,3</sup> and Craig Marston<sup>2,3</sup><sup>1</sup>CSIRO Health & Biosecurity, Tropical Ecosystems Research Centre, PMB 44, Winnellie, NT, 0822, Australia<sup>2</sup>Department of Agriculture, Water and the Environment, 42-44 Qantas Dr, Brisbane, Queensland, 4008, Australia<sup>3</sup>Current address: Department of Agriculture Fisheries and Forestry, 42-22 Qantas Dr, Brisbane, Queensland, 4008, AustraliaORCID: [0000-0002-4010-4723](https://orcid.org/0000-0002-4010-4723) (BDH)Corresponding author: Benjamin D. Hoffmann ([Ben.Hoffmann@csiro.au](mailto:Ben.Hoffmann@csiro.au))

**Citation:** Hoffmann BD, Eldridge J, Marston C (2023) The first eradication of an exotic ant species from the entirety of Australia: *Pheidole fervens*. *Management of Biological Invasions* 14(4): 619–624, <https://doi.org/10.3391/mbi.2023.14.4.03>

**Received:** 30 March 2023**Accepted:** 14 September 2023**Published:** 9 November 2023**Handling editor:** Staci Amburgey**Thematic editor:** Catherine Jarneveich**Copyright:** © Hoffmann et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International - CC BY 4.0).

**OPEN ACCESS****Abstract**

In July 2019 an established infestation of *Pheidole fervens* was detected in Australia for the first time. An eradication program was conducted, and by November 2022 the ant had not been detected for 31 months so it was declared eradicated. This is the first time an exotic ant species has been eradicated from within Australia without other infestations of the same species being found in the meantime. Much of the achievement can be attributed to early detection of the incursion by Australia's National Border Surveillance program.

**Key words:** ants, ecology, Hymenoptera: Formicidae, impacts, invasive**Introduction**

Many ant species have become dispersed throughout the world outside of their native range (McGlynn 1999; Wong et al. 2023), with a great variety of consequences (Holway et al. 2002; Gruber et al. 2022) and associated costs (Angulo et al. 2022), increasingly requiring management action for control or eradication (Hoffmann et al. 2016). The ant species *Pheidole fervens* (Figure 1) is one such species that has been accidentally dispersed to so many places globally, and with much of that dispersal happening so long ago, that there is uncertainty over its origins as native to either SE Asia or the Pacific (Wilson and Taylor 1967; Eguchi 2004; Sarnat et al. 2015). In 2019 an established (i.e. having a viable colony living within local substrate, not associated with the goods within which it was originally transported) infestation of *P. fervens* was detected in Australia for the first time. Although this species is not known to have any serious environmental, social or agricultural impacts, an eradication program commenced based on the precautionary principle that it would be better to eliminate the incursion before any potential impacts became realised, and the species had spread so widely that eradication becomes impossible or prohibitively expensive. Here we describe that eradication, which has now been successfully completed, coincidentally being the first time an exotic ant has



**Figure 1.** *Pheidole fervens* major worker collected from the incursion. Photograph by Tony Robinson.

been eradicated from within Australia without other infestations of the same species being found in the meantime.

### Methods and results

In July 2019, lures were placed at a bulk terminal facility (a Biosecurity Entry Point) at Murrarie, Port of Brisbane, Queensland (22.44921389°S; 153.09516667°E), as part of Australia's National Border Surveillance program which routinely surveys locations around Australia where new incursions of exotic species may establish. Lures were 50 mL urine sample jars containing either protein in the form of a slice of hot dog sausage (Don Footy Franks®), or carbohydrate in the form of a half teaspoon of golden syrup, positioned randomly throughout the area. Lures were placed on the ground on their side with their lid off for approximately one hour before being collected. Subsequent laboratory assessment of the captured ants identified *P. fervens*. Specimens had been found in lures that had been placed on the ground along a concrete gutter structure. Additional specimens were hand collected from the same location soon afterwards and sent to the Okinawa Institute of Science & Technology in Japan for DNA confirmation. Identification was confirmed on 21 August 2019.

The incursion was delimited in September 2019. First, approximately 30 ha was assessed visually, with assessments extending out 300 m from detection locations. Lures were then used to refine the infested areas. Lures with different food sources were alternated on the ground and were spaced in an approximate 5 × 5 m grid within the visually estimated infested areas. An approximate 10 × 10 m grid was used in the surrounding non-infested areas extending out 100 m from the periphery of the visual detection locations. The assessments used approximately 1250 lures, and the ant was found to inhabit a combined



**Figure 2.** Detection points (red points) for *Pheidole fervens* and areas treated by spray (blue polygons) or granular product (orange polygons).

area of 1.84 ha in two spatially discrete areas covering 14,400 m<sup>2</sup>, and 4000 m<sup>2</sup> on one property (Figure 2). The two infestations were separated by approximately 350 m. An eradication plan was subsequently drafted and underwent a rapid consultation process with biosecurity representatives from all Australian States and Territories, as well as ant eradication specialists. In October 2019 the eradication plan was finalised (Department of Agriculture 2019). On 13 October 2019, a spray treatment of Termidor® Residual Termiticide and Insecticide containing the active ingredient fipronil (100 g/L) was applied over six areas containing exposed ground (i.e. not sealed roads) up to 5 m from the river (Figure 2). The spray mixture was prepared as per the product label (BASF 2019) using 6 mL Termidor per 1 L water and applied at the rate of 1 L per 25 lineal metres. The area within 5 m of the river was treated using the granular product Distance® Plus containing the active constituent pyriproxyfen (5 g/kg), applied at the rate of 4 kg/ha (Sumitomo Chemical Australia 2019).

The combined 1.84 ha of infested areas was surveyed again in January 2020 using 525 lures in the same manner as detailed for the delimitation survey (Supplementary material Figure S1), with two individuals of *P. fervens* being found in two adjacent lures. In March 2020, a second treatment was applied. Additionally, green-waste piles approximately 50 m from the infestation were removed, and placed in layers into large industrial bins, with each layer sprayed as they were placed in the bins with Termidor® at the concentration detailed prior.

Additional post treatment assessments were conducted in May and November 2020, and November 2021 throughout the same 1.8 ha, with no *P. fervens* being collected within 321, 1591 and 1339 lures respectively (Figure S1). However, in order to verify distribution had not spread to neighbouring areas another survey was conducted incorporating areas up to 300 m away from the infested areas. Between 26 October 2022 and 24 November 2022, an additional 1304 lures were deployed over approximately 16 ha (Figure S1), with no *P. fervens* being found. The ant was subsequently

declared eradicated, with its actual eradication date likely occurring in March 2020 immediately following the second treatment.

The treatments had a minor and short-term effect on other ant species in the treatment area. Pre-treatment, 27 species (excluding *P. fervens*) were present, including another exotic *Pheidole* the African big-headed ant *P. megacephala* (Table S1). In the three post-treatment assessments of the treated area 14, 25 and 25 species were found respectively, but not *P. megacephala*. In the final assessment over a broader area that wasn't treated, 31 species were found.

## Discussion

There are at least 20 exotic ant species established in Australia (Majer and Heterick 2015) but this is the first established species to have all known infestations eradicated. Despite numerous site-level and regional eradications of other species including yellow crazy ant *Anoplolepis gracilipes*, red imported fire ant *Solenopsis invicta*, tropical fire ant *Solenopsis geminata*, African big-headed ant *Pheidole megacephala*, Argentine ant *Linepithema humile*, browsing ant *Lepisiota frauenfeldi*, and little fire ant *Wasmannia auropunctata* (Hoffmann et al. 2016, 2023; Wylie et al. 2016), all of these species still have infestations remaining extant elsewhere in Australia.

Possibly the greatest reason that *P. fervens* has been the first species eradicated from within Australia without other infestations of the same species being found in the meantime, as well as being done so quickly and for only AUD\$30,000, is that the incursion appears to have been detected at a very early stage, and prior to secondary dispersal events that would have extended work to other locations. Notably, previous surveillance conducted at this location in April 2017 had not detected *P. fervens*, so presumably population levels were so low and small at that point that they were not detected, or the incursion occurred after April 2017. Other rapid eradications that can be attributed to early detections are the three declared eradications of *S. invicta* from New Zealand in 2003, 2007 and 2009 (Pascoe 2003; Sarty 2007; Christian 2009), Port Botany in Sydney in 2017 (New South Wales Government *unpublished data*), Brisbane airport in 2019 (Queensland Government *unpublished data*), and Port of Brisbane in 2021 (Queensland Government *unpublished data*), all of which were single nests when found. All these detections were made within national surveillance programs and illustrate the benefit of conducting such surveillance.

Although the protocols used here successfully achieved eradication, we acknowledge that normally the treatments would have included a buffer area extending away from the infested area to accommodate the potential that the infested area had not been delimited accurately (FAO 1988). We are not able to offer a reason why a buffer zone was not included in the protocols used here, but note that a best-practice document is still lacking for invasive ant eradication protocols for anywhere in the world. Therefore, even within



Australia, which is conducting numerous eradication programs, there is great variability in the protocols being used by different entities focused on the same species. However, we recommend all future programs treat a buffer zone around the infested area. This is the first time an exotic ant species has been eradicated from within Australia without other infestations being found in the meantime. We acknowledge that this species may indeed be currently undetected in another location, but this is much more unlikely than in the past, given that the Australia's National Biosecurity Program now frequently assesses primary and secondary destinations of international goods, and there is also an immense number of locations being assessed simultaneously as part of the numerous ant eradication programs currently occurring within Australia.

As international trade increases globally, and subsequently exotic species are unintentionally and increasingly travelling in cargo (Essl et al. 2011; Seebens et al. 2017), it is highly likely that more incursions of *P. fervens* will occur. It is also possible for this incursion that a secondary infestation has occurred and remains unknown and will be found in the future. In both instances, the work presented here has demonstrated that eradication of such an incursion is achievable, and is no doubt more economically viable than the cost of living with such exotic species (Angulo et al. 2022).

## Acknowledgements

We acknowledge Ramandeep Kaur for first identifying *P. fervens* from the monitoring samples, and with Mizuki Uemura and Andrew Maynard identifying all ants in the laboratory. Sarah Johnston, Jamie Summerhayes, Sarah Hickman, Todd Spencer, Kurt Wadley, and Daniel Contarini conducting the ground work associated with the eradication. Tony Robinson provided the photograph for Figure 1. We thank Marc Widmer, Ross Wylie, and two anonymous reviewers for reviewing the draft manuscript. The eradication program was funded by the Australian Department of Agriculture through the National Border Surveillance Program.

## Authors' contribution

JE and CM led the eradication. BH, JE and CM drafted the manuscript. All authors edited and approved the final manuscript.

## References

- Angulo E, Hoffmann BD, Ballesteros-Mejia L, Taheri A, Balzani P, Renault D, Cordonnier M, Bellard C, Diagne C, Ahmed DA, Watari Y, Courchamp F (2022) Economic costs of invasive alien ants worldwide. *Biological Invasions* 24: 2041–2060, <https://doi.org/10.1007/s10530-022-02791-w>
- Christian S (2009) Red imported fire ants eradicated from Napier. *Biosecurity Magazine* 92: 28–29
- Department of Agriculture (2009) Introduced invasive ant *Pheidole fervens* treatment plan. National Border Surveillance. Department of Agriculture, Brisbane, 12 pp
- Eguchi K (2004) Taxonomic revision of two wide-ranging Asian ants, *Pheidole fervens* and *P. indica* (Insecta: Hymenoptera, Formicidae) and related species. *Annals of the Naturhistorisches Museum Wien* 105B: 189–209
- Essl F, Dullinger S, Rabitsch W, Hulme PE, Hülber K, Jarošík V, Kleinbauer I, Krausmann F, Kühn I, Nentwig W, Vilà, Genovesi P, Gherardi F, Desprez-Loustau ML, Roques A, Pyšek P (2011) Socioeconomic legacy yields an invasion debt. *Proceedings of the National Academy of Sciences* 108: 203–207, <https://doi.org/10.1073/pnas.1011728108>
- Gruber MAM, Santoro D, Cooling M, Lester PJ, Hoffmann BD, Boser C, Lach L (2022) A global review of socio-economic and environmental impacts of ants reveals new insights for risk assessment. *Ecological Applications* 32: e2577, <https://doi.org/10.1002/eap.2577>

- Hoffmann BD, Luque GM, Bellard C, Holmes ND, Donlan CJ (2016) Improving invasive ant eradication as a conservation tool: A review. *Biological Conservation* 198: 37–49, <https://doi.org/10.1016/j.biocon.2016.03.036>
- Hoffmann BD, Widmer M, Bates OK (2023) Preparing to eradicate a novel invader of unknown biology: a case study from Australia. *Management of Biological Invasions* 14: 421–436, <https://doi.org/10.3391/mbi.2023.14.3.03>
- Holway D, Lach L, Suarez A, Tsutsui N, Case T (2002) The causes and consequences of ant invasions. *Annual Review of Ecology, Evolution and Systematics* 33: 181–233, <https://doi.org/10.1146/annurev.ecolsys.33.010802.150444>
- Majer JD, Heterick BE (2015) Invasive ants on the Australian mainland: the other 24 species. Proceedings of the XXII Simpósio de mirmecologia: ann international ant meeting, 18–22 October 2015, Ilhéus, Bahia, Brazil. Centre for Cocoa Research, Brazil
- McGlynn TP (1999) The worldwide transfer of ants: geographical distribution and ecological invasion. *Journal of Biogeography* 26: 535–548, <https://doi.org/10.1046/j.1365-2699.1999.00310.x>
- Pascoe A (2003) Red imported fire ant response stood down. *Biosecurity Magazine* 45: 7
- Sarnat EM, Fischer G, Guenard B, Economo EP (2015) Introduced *Pheidole* of the world: taxonomy, biology and distribution. *Zookeys* 543: 1–109, <https://doi.org/10.3897/zookeys.543.6050>
- Sarty M (2007) Fire ant eradicated at Port of Napier. *Biosecurity* 73: 10
- Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, Winter M, Arianoutsou M, Bacher S, Blasius B, Brundu G, Capinha C, Celesti-Grapo L, Dawson W, Dullinger S, Fuentes B, Jäger H, Kartesz J, Kenis M, Kreft H, Kühn I, Lenzner B, Liebhold A, Mosena A, Moser D, Nishino M, Pearman D, Pergl J, Rabitsch W, Rojas-Sandoval J, Roques A, Rorke S, Rossinelli S, Roy HE, Scalera R, Schindler S, Štajerová K, Tokarska-Guzik B, van Kleunen M, Walker K, Weigelt P, Yamanaka T, Essl F (2017) No saturation in the accumulation of alien species worldwide. *Nature Communications* 8: 14435, <https://doi.org/10.1038/ncomms14435>
- Wilson EO, Taylor RW (1967) The ants of Polynesia (Hymenoptera: Formicidae). *Pacific Insects Monographs* 14: 1–109
- Wong MKL, Economo EP, Guénard B (2023) The global spread and invasion capacities of alien ants. *Current Biology* 33: 566–571, <https://doi.org/10.1016/j.cub.2022.12.020>
- Wylie R, Jennings C, McNaught M, Oakley J, Harris E (2016) Eradication of two incursions of the Red Imported Fire Ant in Queensland, Australia. *Ecological Management & Restoration* 17: 22–32, <https://doi.org/10.1111/emr.12197>

## Websites

- BASF (2019) Termidor® residual termiticide and insecticide product label. [https://pest-control.basf.com.au/sites/basf.com.au/files/2021-01/termidor\\_residual\\_label.pdf](https://pest-control.basf.com.au/sites/basf.com.au/files/2021-01/termidor_residual_label.pdf) (accessed 8 March 2023)
- FAO (1998) Guidelines for pest eradication programmes. International standards for phytosanitary measures no. 9, Rome. Available at <https://www.ippc.int> (accessed 8 March 2023)
- Sumitomo Chemical Australia (2019) Distance® Plus Product Label. <https://sumitomo-chem.com.au/downloads/sds-sln> (accessed 8 March 2023)

## Supplementary material

The following supplementary material is available for this article:

**Figure S1.** Locations of the lure assessments with and without *Pheidole fervens* present before and after treatment.

**Table S1.** Ant species found in pitfall trap samples around the infested premise.

This material is available as part of online article from:

[http://www.reabic.net/journals/mbi/2023/Supplements/MBI\\_2023\\_Hoffmann\\_et\\_al\\_SupplementaryMaterial\\_2.pdf](http://www.reabic.net/journals/mbi/2023/Supplements/MBI_2023_Hoffmann_et_al_SupplementaryMaterial_2.pdf)

Supplementary Material

Recommended citation:


Hoffmann BD, Eldridge J, Marston C (2023) The first eradication of an exotic ant species from the entirety of Australia: *Pheidole fervens*. *Management of Biological Invasions* 14(4): 619–624, <https://doi.org/10.3391/mbi.2023.14.4.03>

**Figure S1.** Locations of the lure assessments with and without *Pheidole fervens* present

**Table S1.** Ant species found in pitfall trap samples around the infested premise




**Figure S1.** Locations of the lure assessments with (red circles) and without (white circles) *Pheidole fervens* present, conducted at A) 3 months after the first treatment and two months before the second treatment in January 2020, B) 2 months post-treatment in May 2020, C) 8 months post-treatment in November 2020, D) 20 months post-treatment in November 2021, and E) 31 months post-treatment in October/November 2022, at the infested premise (22.44921389° S, 153.09516667 E).






C) 8 months post-treatment,  
November 2020



D) 20 months post-treatment,  
November 2021



E) 31 months post-treatment,  
October/November 2022





Table S1. Ant species found in pitfall trap samples around the infested premise (22.44921389° S, 153.09516667 E) within the five sample times: Pre-treatment in July 2019; Post-treatment 1 in May 2020; Post-treatment 2 in November 2020; Post-treatment 3 in November 2021, and a final post-treatment assessment throughout a broader area in October/November 2022.

Species	Pre-treat	Post 1	Post 2	Post 3	Broad area
<i>Brachyponera lutea</i>	X				
<i>Camponotus aeneopilosus</i>	X				
<i>Camponotus gasseri</i>					X
<i>Camponotus humilior</i>	X				
<i>Camponotus sp.</i>			X	X	X
<i>Cardiocondyla atalanta</i>					X
<i>Cardiocondyla emeryi</i>					X
<i>Cardiocondyla nuda</i>	X	X	X	X	X
<i>Crematogaster laeviceps</i>					X
<i>Crematogaster sp.</i>				X	
<i>Doleromyrma sp.</i>	X				
<i>Heteroponera sp.</i>		X	X		
<i>Iridomyrmex fervens</i>					X
<i>Iridomyrmex purpureus</i>			X	X	X
<i>Iridomyrmex rufoniger</i>	X				
<i>Iridomyrmex sp.</i>	X	X	X	X	X
<i>Melophorus sp.</i>				X	
<i>Meranoplus sp.</i>			X	X	X
<i>Monomorium fieldi</i>	X		X	X	X
<i>Monomorium floricola</i>	X	X	X	X	X
<i>Monomorium rothsteini</i>	X				
<i>Monomorium sp.</i>			X	X	X
<i>Notostigma carazzii</i>			X		
<i>Notoncus ectatommoides</i>	X	X			X
<i>Notoncus gilberti</i>	X				
<i>Notoncus sp.</i>			X	X	
<i>Nylanderia braueri</i>	X	X	X	X	X
<i>Nylanderia obscura</i>				X	
<i>Nylanderia sp.</i>				X	X
<i>Ochetellus glaber</i>	X		X	X	X
<i>Opisthopsis rufithorax</i>		X	X	X	
<i>Papyrius nitidus</i>		X			
<i>Paraparatrechina sp.</i>	X				
<i>Paratrechina longicornis</i>	X		X	X	X
<i>Pheidole megacephala</i>	X				X
<i>Pheidole rugulosa</i>		X	X	X	X
<i>Pheidole sp.</i>	X				X

<i>Plagiolepis alluaudi</i>	X		X	X	X
<i>Podomyrma gratiosa</i>					X
<i>Polyrhachis australis</i>	X	X			
<i>Polyrhachis vermiculosa</i>	X				
<i>Polyrhachis</i> sp.			X	X	
<i>Ponera</i> sp.			X		
<i>Rhytidoponera metallica</i>	X	X	X	X	X
<i>Rhytidoponera</i> sp.		X			
<i>Strumigenys</i> sp					X
<i>Tapinoma melanocephalum</i>	X		X	X	
<i>Tapinoma</i> sp.	X	X	X		X
<i>Technomyrmex jocosus</i>	X		X	X	X
<i>Technomyrmex sofiae</i>					X
<i>Tetramorium bicarinatum</i>			X	X	
<i>Tetramorium impressum</i>					X
<i>Tetramorium lanuginosum</i>	X				X
<i>Tetramorium simillimum</i>	X	X	X	X	X