

- GENUNG, W. G., AND M. J. JANES. 1975. Host range, wild host significance, and in-field spread of *Liriomyza trifolii* and population build-up and effects of its parasites in relation to Fall and Winter celery (Diptera: Agromyzidae). Belle Glade AREC Res. Rpt. EV-1975-5, 18 p.
- GILSTRAP, F. E., AND E. R. OATMAN. 1976. The bionomics of *Scolothrips sexmaculatus* (Pergande) (Thysanoptera: Thripidae) an insect predator of spider mites. Hilgardia 44: 27-59.
- LEIBEE, G. L. 1981. Insecticidal control of *Liriomyza* spp. on vegetables, pp. 216-220. in D. J. Schuster [ed.], Proceedings of the IFAS-Ind. Conference on the Biology and Control of *Liriomyza* Leafminers. IFAS, Univ. of Florida.
- NUESSLY, G. S., AND R. T. NAGATA. 1993. Evaluation of damage by serpentine leafminer and banded cucumber beetles to cos lettuce. Everglades Res. and Ed. Center Res. Rpt., EV-1993-2:76-77.
- NUESSLY, G. S., AND R. T. NAGATA. 1994. Differential probing response of serpentine leafminer, *Liriomyza trifolii* (Burgess), on cos lettuce. J. Entomol. Sci. 29: 330-338.
- PARRELLA, M. P., AND K. L. ROBB. 1982. Technique for staining eggs of *Liriomyza trifolii* within chrysanthemum, celery, and tomato leaves. J. Econ. Entomol. 75: 383-384.
- SIMONET, E. E., AND R. L. PIENKOWSKI. 1977. Sampling and distribution of potato leafhopper eggs in alfalfa stems. Ann. Entomol. Soc. America 51: 557-566.
- SUGIMOTO, T. 1976. On distribution of egg population of a leafmining fly, *Phytomyza ranunculi* Schrank (Diptera, Agromyzidae) among leaves and in a leaf. Fac. Agric. Kinki Univ. 9: 11-19.



COMMUNITY STRUCTURE OF HOUSE-INFESTING ANTS  
(HYMENOPTERA: FORMICIDAE) IN SOUTHERN BAHIA,  
BRAZIL

JACQUES H. C. DELABIE<sup>1,3</sup>, IVAN C. DO NASCIMENTO<sup>1,3</sup>, PRISCILA PACHECO<sup>2,3</sup>  
AND ANTONIO B. CASIMIRO<sup>1,3</sup>

<sup>2</sup>Departamento de Enfermagem,

Universidade Estadual de Santa Cruz (UESC), 45660,  
Ilhéus, Bahia, Brazil.

<sup>3</sup>Laboratório de Mirmecologia, Centro de Pesquisas do Cacau (CEPEC),  
CEPLAC, C.P. 7, 45600,  
Itabuna, Bahia, Brazil.

## ABSTRACT

The community structure of ants which infest houses in southern Bahia was studied. Of the 31 species collected, 8 are considered as characteristic of human habitat, even though some of them are also found in other conditions, such as in regional agro-systems. *Pheidole megacephala* was the dominant species and showed a negative association with most other ants, while *Tapinoma melanocephalum* was almost always found with other ant species. The notable absence of the species *L. humile* in our tests may be due to the antagonistic effects of *Ph. megacephala*.

Key Words: *Tapinoma melanocephalum*, *Pheidole megacephala*, *Linepithema humile*, Bahia, Brazil.

#### RESUME

La structure de la communauté de fourmis qui infestent les habitations dans le sud de Bahia a été étudiée. Parmi les 31 espèces rencontrées, 8 sont considérées caractéristiques de l'habitat humain, bien qu'elles soient aussi rencontrées dans d'autres conditions, telles que dans certains agrosystèmes régionaux. Dans les communautés, *Pheidole megacephala* est dominante et montre une association négative avec la plupart des autres fourmis, alors que *Tapinoma melanocephalum* est à peu près toujours rencontrée avec d'autres espèces. L'absence notable de *Linepithema humile* de nos tests est probablement liée à son antagonisme avec *Ph. megacephala*.

---

Ants which infest houses have only recently received attention from entomologists in South America (Brown 1964, Fowler et al. 1992, 1993, Ketelhut et al. 1993; Bueno & Fowler 1994) in contrast to the Northern hemisphere, where ants which occur in houses and hospitals have been studied for at least two or three decades (see the reviews of Smith 1965, Edwards 1986, Eichler 1990, Thompson 1990). Compared to temperate environments, urban tropical environments may yield ideal conditions for a range of species for foraging and nesting. However, in both the temperate and tropical urban conditions of Americas, ants which occur in the human environment are regarded as destructive or detrimental to human health, food conservation and quality, wood conservation, electrical installations, and electronic equipment (Smith 1965, Fowler 1990, Thompson 1990, Vinson & McKay 1990, Fowler et al. 1993a & b).

A higher frequency of occurrence of ants in houses in tropical regions of South America compared to houses in temperate regions (Fowler et al. 1993, Bueno & Fowler 1994) is expected because of the favorable climatic conditions and the characteristics of human habitation in the tropics. To understand the community structure of house-infesting ants in tropical regions, a cooperative study was conducted in the State of Bahia, Brazil, by the Universidade Estadual de Santa Cruz (UESC) and the Myrmecology Laboratory of the Cocoa Research Center (CEPEC), at Ilheus, with the aim of describing the ant situation in houses of this region. This study is the first step toward a larger study on the interactions between ants and human health in tropical environments.

#### MATERIAL AND METHODS

One hundred houses were randomly sampled for ants in the region of Ilheus (14°45'S, 39°13'W), in the southern portion of the State of Bahia. Ants were caught in small glass test-tubes (length: 50 mm; diam: 7 mm) using honey as bait. In each house, three test-tubes were placed on the floor in each room: livingroom, bedroom, kitchen, and bathroom. Tubes were collected in the morning, 8 to 12 hours after initial placement. Collected test-tubes were transported to the Myrmecology Laboratory of the CEPEC where the ants were identified to species.

Data were analysed using a house or a room as the study unit. An evaluation of interrelations between the most common house-infesting ant species and their possible associations with other species were tested by chi-square analysis with Yates' correction applied (Siegel 1956). This procedure has previously been used by Room (1971) and Majer et al. (1994) to study the ant mosaic in cocoa plantations.




Fig. 1. Frequency of houses infested by ants as a function of the number of species found per house.

#### RESULTS AND DISCUSSION

All houses sampled were occupied by at least one species of ant. A total of 31 species belonging to 14 genera and 4 subfamilies were found: Ponerinae: *Gnamptogenys* [1 sp.], *Odontomachus* [1 sp.]; Myrmicinae: *Acromyrmex* [1 sp.], *Crematogaster* [1 sp.], *Monomorium* [2 spp.], *Pheidole* [5 spp.], *Solenopsis* [4 spp.], *Tetramorium* [3 spp.], *Wasmannia* [1 sp.]; Dolichoderinae: *Dorymyrmex* [1 sp.], *Tapinoma* [1 sp.] and Formicinae: *Brachymyrmex* [1 sp.], *Camponotus* [7 spp.], *Paratrechina* [2 spp.]. Eight exotic species (25.8%), *Monomorium floricola* (Jerdon), *M. pharaonis* (L.), *Paratrechina longicornis* (Latr.), *Pheidole megacephala* (Fabr.), *Tapinoma melanocephalum* (Fabr.), *Tetramorium bicarinatum* (Nyl.), *T. lucayanum* Wheeler and *T. simillimum* (Fr. Smith) (Delabie 1993) were responsible for 68.9% of the total number of occurrences. Most of these ants are known as "tramp ants" and have characteristics in common, such as polygyny, low intraspecific aggressivity, and the ability to change their nest site easily (Passera 1993).

One to six species were found per house, and the frequency observed for each of these six classes decreased inversely to the number of ants found (Fig. 1). The greatest number of ant occurrences (29%) and species (22) was observed in the livingroom (Fig. 2A,B). In Bahia, the livingroom is generally the largest room in the house and probably also offers more possibilities for nest sites (e.g., plant containers) to a variety of species. The other three rooms sampled (bathroom, bedroom and kitchen) showed a similar range of species number (14-16) and ant species occurrence (22-27%) (Fig. 2A,B). The kitchen was the location with the highest number of species and occurrences and the bedroom the lowest.

Among the 31 species collected, eight (Table 1) could be considered to be truly characteristic of human habitats because they were found at different sites and in all parts

TABLE 1. RELATIVE FREQUENCY OF MOST COMMON SPECIES OF HOUSE-INFESTING ANTS IN SOUTHERN BAHIA (&gt;10 OCCURRENCES AND LIVING EVERYWHERE IN HOUSES).

| Ant species                        | % of Houses | % of Occurrences |
|------------------------------------|-------------|------------------|
| <i>Pheidole megacephala</i>        | 47.0        | 31.7             |
| <i>Tapinoma melanocephalum</i>     | 48.0        | 22.9             |
| <i>Solenopsis saevissima</i>       | 23.0        | 9.3              |
| <i>Paratrechina longicornis</i>    | 18.0        | 7.9              |
| <i>Wasmannia auropunctata</i>      | 12.0        | 5.5              |
| <i>Camponotus (Tanaemyrmex) sp</i> | 13.0        | 4.8              |
| <i>Pheidole sp4</i>                | 13.0        | 3.8              |
| <i>Tetramorium simillimum</i>      | 9.0         | 2.9              |

of the house. These eight species were responsible for 88.9% of the total number of occurrences. In Bahia, these ants are not found exclusively in houses, but are also commonly found in gardens, secondary vegetation, and crops. Their distribution tends to be different at higher latitudes where few species are limited exclusively to human dwellings, e.g., *M. pharaonis* (Eichler, 1990). This difference is probably related to the year-round thermal stability in this region, which allows these ants to live both in and out of human habitats.

The most abundant ant species were *Ph. megacephala* (47% of infested houses and 31.7% of occurrences) and *T. melanocephalum* (48% of infested houses and 22.9% of occurrences).

The community structure at the room level (Fig. 3) showed that *Ph. megacephala* was the only species that was truly dominant (see Majer et al. 1994). This ant was negatively associated with the species *P. longicornis*, *Solenopsis saevissima* (Fab.), *T. melanocephalum* and the little fire ant *Wasmannia auropunctata* (Roger). *Ph. megacephala* was frequently the only species in a room or a house, with all potential competitors being excluded (Fig. 4). In contrast, the second most common species, *T. melanocephalum*, usually occurred with other ants (Fig. 4). However, it seems that




Fig. 2. Percent of ant occurrences (A) and number of species (B) in sampled rooms.




Fig. 3. Community structure of house-infesting ants of Bahia.

this species can act as a rapid colonizer, because it was the only species found when houses were new.

The study of size classes shows a bimodal distribution (Fig. 5) in which the smaller ants (< 5mm) were more common. Most of the larger ants were species of the *Camponotus* genus which are generally more active at night in Bahia (see, for example, Delabie et al. 1991). Establishment of these large ant species in the house environment may relate to their nocturnal behavior which does not interfere with the human activity and removes them from competition with diurnal species.




Fig. 4. Relative abundance of the 5 commonest ant species in Bahia houses.




Fig. 5. Distribution of house-infesting ants as a function of body length.

*Linepithema humile* (Mayr) was completely absent from our house samples. This absence has been confirmed in all of south Bahia and confirmed by other extensive experiments over the region (J. H. C. D., unpublished data). However, the same species was commonly found forming discrete societies in cocoa plantations within the same region. It seems that *L. humile* avoids human settings in Bahia and this may be due to an antagonism with *Ph. megacephala* which is dominant in houses and other strongly human anthropized habitats. The antagonism between these two species has been documented in other situations (Haskins & Haskins 1965, Crowell 1968). Thus, the main difference between house-infesting ant communities in Bahia and other parts of Brazil is that *L. humile* can be found everywhere in the urban environment (Brown 1964, Fowler et al. 1993), while it seems restricted to only a few agricultural situations in Bahia.

#### ACKNOWLEDGMENTS

We wish to thank Dr. David F. Williams and three reviewers for valuable comments and corrections on an earlier version of this paper. This research was granted by the Universidade Estadual de Santa Cruz (project "House-infesting ants and human health"). Paper presented at the IV International Symposium on Pest Ants, Belo Horizonte (Brazil), November 1993.

#### REFERENCES CITED

- BROWN, W. L., JR. 1964. Some tramp ants of Old World origin collected in Tropical Brazil. *Entomol. News* 75: 14-15.  
 BUENO, O. C., AND H. G. FOWLER. 1994. Exotic ants and the ant fauna of Brazilian hospitals, pp. 191-198 in D. F. Williams [ed.] *Exotic ants: biology, impact, and control of introduced species*, Westview Press, Boulder.

- CROWELL, K. L. 1968. Rates of competitive exclusion by the Argentine ant in Bermuda. *Ecology* 49: 551-555.
- DELABIE, J. H. C. 1993. Formigas exóticas na Bahia. Bahia, Análise e Dados, CEI, 3: 19-22.
- DELABIE, J. H. C., F. P. BENTON, AND M. A. DE MEDEIROS. 1991. La polydomie chez les Formicidae arboricoles dans les cacaoyères du Brésil: optimisation de l'occupation de l'espace ou stratégie défensive? *Actes Coll. Insectes Sociaux* 7: 173-178.
- EDWARDS, J. P. 1986. The biology, economic importance and control of the pharaoh's ant, *Monomorium pharaonis* L., pp. 257-271 in S. B. Vinson [ed.], *Economic impact and control of social insects*. Praeger Press, New York.
- EICHLER, W. 1990. Health aspects and control of *Monomorium pharaonis*, pp. 671-675 in R. K. Vander Meer, K. Jaffe and A. Cedeno [eds], *Applied Myrmecology: a World Perspective*, Westview Press, Boulder.
- FOWLER, H. G. 1990. Carpenter ants (*Camponotus* spp): pest status and human perception, pp. 525-532 in R. K. Vander Meer, K. Jaffe and A. Cedeno [eds], *Applied Myrmecology: a World Perspective*, Westview Press, Boulder.
- FOWLER, H. G., F. ANARUMA FILHO, AND O. C. BUENO. 1992. Vertical and horizontal foraging: intra and interspecific spatial autocorrelation patterns in *Tapinoma melanocephalum* and *Monomorium pharaonis* (Hymenoptera: Formicidae). *Ciência e Cultura* 44: 395-397.
- FOWLER, H. G., F. ANARUMA FILHO, AND O. C. BUENO. 1993a. Seasonal space usage by the introduced pharaoh's ant, *Monomorium pharaonis* (L.) (Hym., Formicidae), in institutional settings in Brazil and its relation to other structural ant species. *J. Appl. Entomol.* 115: 416-419.
- FOWLER, H. G., O. C. BUENO, T. SADATSUNE, AND A. C. MONTELLI. 1993b. Ants as potential vectors of pathogens in hospitals in the State of São Paulo, Brazil. *Insect Sci. Applic.* 14: 367-370.
- HASKINS, C. P., AND E. F. HASKINS. 1965. *Pheidole megacephala* and *Iridomyrmex humilis* in Bermuda - equilibrium or slow displacement? *Ecology* 46: 736-740.
- KETELHUT, S. M., H. G. FOWLER, M. D. GUEDES, B. INDJAI, M. CASTELLANI, AND O. C. BUENO. 1993. Formigas caseiras de Rio Claro-SP (Hymenoptera: Formicidae). *Resumos, IV International Symposium on Pest Ants*, Belo Horizonte, Brazil.
- MAJER, J. D., J. H. C. DELABIE, AND M. R. B. SMITH. 1994. Arboreal ant community patterns in Brazilian cocoa farms. *Biotropica* 26: 73-83.
- PASSERA, L. 1993. Quels sont les caractères étho-physiologiques des "fourmis vagabondes"? *Actes Coll. Insectes Sociaux*, 8: 39-45.
- ROOM, P. M. 1971. The relative distribution of ant species in Ghana's cocoa farms. *J. Anim. Ecol.* 40: 735-751.
- SIEGEL, S. 1956. *Nonparametric Statistics for Behavioral Sciences* (Estatística Não-paramétrica para as Ciências do Comportamento - tradução, 1975), McGraw-Hill, São Paulo, 350 pp.
- SMITH, M. R. 1965. House-infesting ants of the Eastern United States, their recognition, biology and economic importance. *Agricultural Research Service, U.S.D.A., Tech. Bull.* 1326, Washington, 105 pp.
- THOMPSON, C. R. 1990. Ants that have pest status in the United States, pp. 51-67 in R. K. Vander Meer, K. Jaffe and A. Cedeno [eds], *Applied Myrmecology: a World Perspective*, Westview Press, Boulder.
- VINSON, S. B. AND W. P. MCKAY. 1990. Effects of the fire ant, *Solenopsis invicta*, on electrical circuits and equipment, pp. 496-503 in R. K. Vander Meer, K. Jaffe and A. Cedeno [eds], *Applied Myrmecology: a World Perspective*, Westview Press, Boulder.

## EFFECTS OF FOREST MANAGEMENT PRACTICES ON TERRESTRIAL COLEOPTERAN ASSEMBLAGES IN SAND PINE SCRUB

CATHRYN H. GREENBERG<sup>1</sup> AND MICHAEL C. THOMAS<sup>2</sup>

<sup>1</sup>USDA Forest Service Southeastern Forest Experiment Station  
and University of Florida, Department of Wildlife and Range Sciences  
118 Newins-Zeigler Hall  
P.O. Box 14524  
Gainesville, Florida 32604

<sup>2</sup>Florida Department of Agriculture and Consumer Services  
Division of Plant Industry  
Florida State Collection of Arthropods  
P.O. Box 147100  
Gainesville, FL 32614-7100

### ABSTRACT

Coleopteran assemblages were sampled monthly for one year using pitfall traps in replicated sites of three 5- to 7-year-old disturbance treatments and mature forested sand pine scrub in the Ocala National Forest, Marion County, Florida. Disturbance treatments were (1) burning at high-intensity and salvage-logging; (2) clearcutting, roller-chopping and broadcast seeding, and; (3) clearcutting and bracke-seeding. Community similarity of coleopterans was high. No differences in species richness, diversity, density, or evenness were detected. Of 40 species captured, only seven were common ( $n > 50$ ). Predaceous beetles were numerically dominant followed by scavengers. Few xylophagous or herbivorous coleopterans were captured, probably due to trap bias. Peaks of annual above-ground terrestrial activity varied among species. An absence of differences among treatments may reflect similar plant communities or structural habitat features. Additionally, a dearth of mature forest specialists might be predicted in systems where mature forest was historically rare due to large-scale, high-intensity, and low-frequency wildfire.

Key Words: Beetle assemblage, clearcutting, wildfire.

### RESUMEN

Durante un año fueron muestreadas mensualmente comunidades de coleópteros usando trampas de suelo en sitios replicados que poseen tres tipos de tratamientos con 5 a 7 años de edad, y el matorral que crece bajo un bosque maduro de pino de arena (sand pine) en el Bosque Nacional de Ocala, en el condado de Marion, en la Florida. Los tratamientos fueron (1) quema de gran intensidad y preservación de los troncos; (2) tala, corte de los troncos en pedazos y siembra de semillas al voleo, y (3) tala y siembra de semillas en los claros talados. La similitud de comunidades de coleópteros fue elevada. No se encontraron diferencias en riqueza de especies, diversidad, o densidad. De 40 especies capturadas solamente siete fueron comunes ( $n > 50$ ). Los escarabajos depredadores fueron numéricamente dominantes, seguidos por los comedores de carroña. Fueron capturados pocos coleópteros xilófagos o herbívoros, probablemente debido a los tipos de trampas. Los picos de actividad terrestre por encima del suelo variaron entre las especies. La falta de diferencia entre los tratamientos puede reflejar comunidades de plantas o formas de habitats estructurales similares. Adicionalmente, la falta de especialistas de bosque maduro podría ser pre-