

NOTOSTIGMA

carazzii. *Camponotus carazzii* Emery, 1895g: 354 (w.) AUSTRALIA (Queensland).

Forel, 1915b: 104 (s.m.); Wheeler, G.C. & Wheeler, J. 1988: 355 (l.).

Combination in *C. (Myrmosphincta)*: Forel, 1912i: 92;

combination in *Notostigma*: Emery, 1920b: 253.

Status as species: Emery, 1896d: 373 (in list); Forel, 1915b: 104; Emery, 1920b: 253; Emery, 1925b: 57; Taylor & Brown, 1985: 127; Taylor, 1987a: 48; Taylor, 1992a: 62; Bolton, 1995b: 293.

Senior synonym of *podenzanai*: Taylor, 1992a: 62; Bolton, 1995b: 293.

foreli. *Notostigma foreli* Emery, 1920b: 253 (w.q.m.) AUSTRALIA (New South Wales).

Status as species: Emery, 1925b: 57; Taylor & Brown, 1985: 127; Taylor, 1987a: 48; Taylor, 1992a: 62; Bolton, 1995b: 293.

podenzanai. *Camponotus podenzanai* Emery, 1895g: 355 (w.m.) AUSTRALIA (Queensland).

Combination in *C. (Myrmosphincta)*: Forel, 1914a: 273;

combination in *Notostigma*: Emery, 1920b: 254.

Status as species: Emery, 1896d: 373 (in list); Forel, 1915b: 105; Emery, 1920b: 254; Emery, 1925b: 57; Taylor & Brown, 1985: 127; Taylor, 1987a: 48.

Junior synonym of *carazzii*: Taylor, 1992a: 62; Bolton, 1995b: 293.

sanguinea Clark, 1930; see under **CAMPONOTUS**.