

LINEPITHEMA

anathema. *Linepithema anathema* Wild, 2007a: 27, figs. 33, 34, 106 (w.) BRAZIL (Minas Gerais, Paraná).

angulatum. *Iridomyrmex humilis* subsp. *angulatus* Emery, 1894c: 165, pl. 1, fig. 3 (w.q.) BOLIVIA.

Wild, 2007a: 31 (m.).

Combination in *Linepithema*: Shattuck, 1992a: 16.

Subspecies of humile: Forel, 1908c: 394; Emery, 1913a: 26; Santschi, 1916e: 390; Luederwaldt, 1918: 47; Santschi, 1923d: 68; Borgmeier, 1923: 87; Santschi, 1929d: 306; Santschi, 1931e: 281; Santschi, 1936d: 417; Kempf, 1972a: 125; Shattuck, 1994: 123; Bolton, 1995b: 246.

Status as species: Wild, 2007a: 29 (redescription); Bezděčková, *et al.* 2015: 109; Escárraga & Guerrero, 2016: 449; Guerrero, 2019: 713.

Senior synonym of pordescens: Wild, 2007a: 29.

arrogans. *Iridomyrmex humilis* var. *arrogans* Chopard, 1921: 237 (footnote), 241, figs. 1-31 (w.q.m.l.) FRANCE.

[Also described as new by Santschi, 1929d: 306 (in key).]

Combination in *Linepithema*: Shattuck, 1992a: 16.

Subspecies of *humile*: Kempf, 1972a: 124; Shattuck, 1992a: 16; Shattuck, 1994: 123; Bolton, 1995b: 246.

Junior synonym of *humile*: Bernard, 1967: 251; Wild, 2004: 1207; Wild, 2007a: 61; Casevitz-Weulersse & Galkowsky, 2009: 478; Radchenko, 2016: 84.

aspidocoptus. *Iridomyrmex aspidocoptus* Kempf, 1969: 295, figs. 10, 11 (w.) BRAZIL (São Paulo).

Combination in *Linepithema*: Shattuck, 1992a: 16.

Status as species: Kempf, 1972a: 124; Shattuck, 1994: 122; Bolton, 1995b: 247.

Junior synonym of *leucomelas*: Wild, 2007a: 81.

aztecoides. *Linepithema aztecoides* Wild, 2007a: 34, figs. 43, 44, 105, 110 (w.) PARAGUAY, BRAZIL (Goiás, Mato Grosso, São Paulo).

Status as species: Wild, 2007b: 24.

bicolor. *Iridomyrmex iniquus* var. *bicolor* Forel, 1912h: 47 (w.) MARTINIQUE I., BRAZIL (Rio de Janeiro)

Combination in *Linepithema*: Shattuck, 1992a: 16.

Subspecies of *iniquum*: Borgmeier, 1923: 88; Kempf, 1972a: 125; Shattuck, 1994: 125; Bolton, 1995b: 247.

Junior synonym of *iniquum*: Wild, 2007a: 68.

breviscapa. *Iridomyrmex humilis* st. *breviscapa* Santschi, 1929d: 306.

[First available use of *Iridomyrmex humilis* r. *platensis* var. *breviscapa* Forel, 1914d: 286 (w.) ARGENTINA (Tucumán); unavailable (infrasubspecific) name.]

Santschi, 1916e: 390 (m.).

Combination in *Linepithema*: Shattuck, 1992a: 16.

As unavailable (infrasubspecific) name: Forel, 1915c: 359.

Subspecies of *humile*: Kempf, 1972a: 125; Zolessi, *et al.* 1988: 7; Shattuck, 1994: 123; Bolton, 1995b: 247.

Junior synonym of *gallardoi*: Wild, 2007a: 55.

bruchi. *Linepithema bruchi* Brandão, 1991: 354. *Nomen nudum.*

- cerradense.** *Linepithema cerradense* Wild, 2007a: 37, figs. 27, 28, 66, 104 (w.q.m.) PARAGUAY, BOLIVIA, BRAZIL (Bahia, Distrito Federal, Goiás, Mato Grosso do Sul, São Paulo).
Status as species: Wild, 2007b: 24.
- cryptobioticum.** *Linepithema cryptobioticum* Wild, 2007a: 41, figs. 9, 10, 100 (w.) PARAGUAY.
Status as species: Wild, 2007b: 24.
- dispertitum.** *Iridomyrmex dispertitus* Forel, 1885a: 351 (w.m.) GUATEMALA. Wild, 2007a: 45 (m.).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Dalla Torre, 1893: 168; Forel, 1899c: 102; Forel, 1908c: 395; Forel, 1912h: 47; Emery, 1913a: 26; Wheeler, W.M. 1916d: 330; Wheeler, W.M. 1922c: 14; Menozzi, 1927c: 268; Kempf, 1972a: 124; Shattuck, 1994: 122; Bolton, 1995b: 247; Wild, 2007a: 43 (redescription); Branstetter & Sáenz, 2012: 254; Escárraga & Guerrero, 2016: 449; Guerrero, 2019: 714; Lubertazzi, 2019: 124.
- dominicensis.** *Iridomyrmex melleus* var. *dominicensis* Wheeler, W.M. 1913d: 242 (w.) DOMINICAN REPUBLIC.
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *melleus*: Kempf, 1972a: 126; Shattuck, 1994: 126; Bolton, 1995b: 247.
Junior synonym of *iniquum*: Wild, 2007a: 68.
- flavescens.** *Iridomyrmex keiteli* var. *flavescens* Wheeler, W.M. & Mann, 1914: 43 (w.) HAITI.
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of keiteli: Wheeler, W.M. 1936b: 204; Kempf, 1972a: 125; Shattuck, 1994: 125; Bolton, 1995b: 247.
Status as species: Wild, 2007a: 49 (redescription); Lubertazzi, 2019: 125.
- fuscescens.** *Iridomyrmex melleus* var. *fuscescens* Wheeler, W.M. 1908a: 153 (w.) PUERTO RICO.
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *melleus*: Emery, 1913a: 27; Smith, M.R. 1937: 864; Kempf, 1972a: 126; Shattuck, 1994: 126; Bolton, 1995b: 247.
Junior synonym of *iniquum*: Wild, 2007a: 68.
- fuscum.** *Linepithema fuscum* Mayr, 1866a: 497, pl. fig. 7 (m.) PERU.
Wild, 2007a: 51 (w.q.).
Status as species: Dalla Torre, 1893: 170; Emery, 1913a: 15; Kempf, 1972a: 135; Shattuck, 1994: 122; Bolton, 1995b: 247; Wild, 2007a: 51 (redescription); Bezděčková, *et al.* 2015: 109.
- gallardoii.** *Dorymyrmex gallardoii* Brèthes, 1914a: 95 (w.) ARGENTINA (Córdoba).
Santschi, 1929d: 306 (q.). Wild, 2007a: 57 (m.).
Combination in *Iridomyrmex*: Gallardo, 1916a: 105;
combination in *Linepithema*: Shattuck, 1992a: 16.
As unavailable (infrasubspecific) name: Santschi, 1931e: 281; Kempf, 1972a: 125.
Subspecies of humile: Gallardo, 1916a: 105; Santschi, 1929d: 306; Shattuck, 1994: 124; Bolton, 1995b: 247.
Status as species: Bruch, 1915: 533; Wild, 2007a: 55 (redescription); Wild, 2007b: 25; Bezděčková, *et al.* 2015: 109; Escárraga & Guerrero, 2016: 450; Guerrero, 2019: 714.
Senior synonym of breviscapa: Wild, 2007a: 55.

Senior synonym of *impotens*: Wild, 2007a: 55.

gallardoi. *Linepithema gallardoi* Kusnezov, 1969: 37 (m.) ARGENTINA (Salta).

[**Junior secondary homonym of *Dorymyrmex gallardoi* Brèthes, 1914a: 95.**]

Status as species: Brandão, 1991: 354; Shattuck, 1994: 122.

Replacement name: *Linepithema inacatum* Bolton, 1995b: 247.

***hirsutum*.** *Linepithema hirsutum* Escárraga & Guerrero, 2016: 450, figs. 1A, 1B, 3 (w.) COLOMBIA.

Status as species: Guerrero, 2019: 714.

***humile*.** *Hypoclinea humilis* Mayr, 1868a: 164 (w.) ARGENTINA (Buenos Aires).

Forel, 1908c: 395 (m.); Newell, 1908: 28 (q.); Wheeler, G.C. & Wheeler, J. 1951: 186 (l.); Crozier, 1969: 250 (k.).

Combination in *H. (Iridomyrmex)*: Mayr, 1870b: 959;

combination in *Iridomyrmex*: Emery, 1888d: 386;

combination in *Linepithema*: Shattuck, 1992a: 16.

Status as species: Mayr, 1870b: 959 (in key); Emery, 1888c: 362; Dalla Torre, 1893: 169; von Jhering, 1894: 378; Forel, 1895b: 110; Forel, 1901h: 81; Emery, 1906c: 177; Forel, 1908c: 394; Wheeler, W.M. 1910g: 569; Santschi, 1912e: 533; Forel, 1912h: 46; Emery, 1913a: 26; Forel, 1913j: 223; Forel, 1913l: 242; Forel, 1914e: 12; Bruch, 1914: 226; Arnold, 1915: 145 (redescription); Bruch, 1915: 534; Donisthorpe, 1915d: 342; Gallardo, 1915: 23; Gallardo, 1916a: 97; Santschi, 1916e: 390; Stitz, 1917: 347; Luederwaldt, 1918: 47; Wheeler, W.M. 1922a: 921; Borgmeier, 1923: 87; Stärcke, 1926: 117 (in key); Essig, 1926: 864; Donisthorpe, 1927a: 7; Donisthorpe, 1927b: 394; Wheeler, W.M. 1927g: 113; Menozzi, 1929a: 4; Wheeler, W.M. 1932a: 12; Santschi, 1933a: 22; Santschi, 1934d: 280; Eidmann, 1936b: 88; Novák & Sadil, 1941: 97 (in key); Grandi, 1935: 102; Donisthorpe, 1950a: 339; Creighton, 1950a: 341; Schmitz, 1950: 14; Smith, M.R. 1951a: 836; Kusnezov, 1953b: 339; Wellenius, 1955: 12; Bernard, 1956b: 260; Ceballos, 1956: 309; Smith, M.R. 1958c: 140; Baroni Urbani, 1964b: 57; Bernard, 1967: 251 (redescription); Smith, M.R. 1967: 364; Wilson & Taylor, 1967: 79; Yarrow, 1967: 29; Baroni Urbani, 1968b: 474; Collingwood & Yarrow, 1969: 75; Kempf, 1970a: 26; Baroni Urbani, 1971c: 166; Kempf, 1972a: 124; Baroni Urbani, 1974: 236; Bolton & Collingwood, 1975: 7 (in key); Hunt & Snelling, 1975: 22; Pisarski, 1975: 27; Snelling, R.R. & Hunt, 1976: 84; Arnol'di & Dlussky, 1978: 548 (in key); Báez & Ortega, 1978: 189; Collingwood, 1978: 87 (in key); Collingwood, 1979: 33; Smith, D.R. 1979: 1418; Snelling, R.R. & George, 1979: 160; Barquin Diez, 1981: 343; Schembri & Collingwood, 1981: 420; Allred, 1982: 478; Wheeler, G.C. & Wheeler, J. 1986g: 56; Agosti & Collingwood, 1987b: 278 (in key); Taylor, 1987a: 32; Ward, 1987: 1; Zolessi, *et al.* 1988: 7; Deyrup, *et al.* 1989: 99; Wheeler, G.C. & Wheeler, J. 1990a: 465; Hohmann, *et al.* 1993: 159; Shattuck, 1994: 123; Douwes, 1995: 91; Bolton, 1995b: 247; Mei, 1995: 767; Poldi, *et al.* 1995: 6; Collingwood, Tigar & Agosti, 1997: 509; Espadaler, 1997b: 27; Collingwood & Prince, 1998: 20 (in key); Deyrup, *et al.* 2000: 300; Czechowski, *et al.* 2002: 16; Mackay & Mackay, 2002: 260; Deyrup, 2003: 45; Imai, *et al.* 2003: 83; Wetterer & Wetterer, 2004: 215; Wild, 2004: 1207 (redescription); Coover, 2005: 106; MacGown & Forster, 2005: 64; Radchenko, 2005b: 155; Ward, 2005: 26; Cagniant, 2006a: 194; Gómez & Espadler, 2006: 227; Petrov, 2006: 102 (in key); Seifert, 2007: 254; Don, 2007: 161; Wetterer, Espadaler, Ashmole, *et al.* 2007: 31; Wetterer, *et al.* 2007: 12; Wild, 2007a: 61 (redescription); Wild, 2007b: 25; Heterick, 2009: 57; Casevitz-

Weulersse & Galkowsky, 2009: 478; Wetterer, Wild, *et al.* 2009: 187; Solis, *et al.* 2010: 19; Lapeva-Gjonova, *et al.* 2010: 33; Boer, 2010: 14; Collingwood, *et al.* 2011: 415; Borowiec, L. & Salata, 2012: 510; Czechowski, *et al.* 2012: 70; Borowiec, L. & Salata, 2013: 358; Borowiec, L. 2014: 100; Bezděčková, *et al.* 2015: 109; Escárraga & Guerrero, 2016: 451; Lebas, *et al.* 2016: 112; Radchenko, 2016: 84; Deyrup, 2017: 172; Salata & Borowiec, 2018c: 46; Seifert, 2018: 241; Dekoninck, *et al.* 2019: 1154; Guerrero, 2019: 714.

Senior synonym of *arrogans*: Bernard, 1967: 251; Wild, 2004: 1207; Wild, 2007a: 61; Casevitz-Weulersse & Galkowsky, 2009: 478; Radchenko, 2016: 84.

Senior synonym of *riograndensis*: Wild, 2004: 1207; Wild, 2007a: 61; Radchenko, 2016: 84.

impotens. *Iridomyrmex impotens* Santschi, 1923d: 67 (w.) BRAZIL (Santa Catarina).

Combination in *Linepithema*: Shattuck, 1992a: 16.

Status as species: Kempf, 1972a: 125; Shattuck, 1994: 124; Bolton, 1995b: 247.

Junior synonym of *gallardoi*: Wild, 2007a: 55.

inacatum. *Linepithema inacatum* Bolton, 1995b: 247.

Replacement name for *Linepithema gallardoi* Kusnezov, 1969: 37. [Junior secondary homonym of *Dorymyrmex gallardoi* Brèthes, 1914a: 95.]

Incertae sedis in *Linepithema*: Wild, 2007a: 18.

iniquum. *Hypoclinea iniqua* Mayr, 1870a: 392, fig. 10 (w.) COLOMBIA (“New Granada”).

Forel, 1908c: 395 (m.); Wild, 2007a: 70 (q.).

Combination in *Iridomyrmex*: Emery, 1890b: 56;

combination in *Linepithema*: Shattuck, 1992a: 16.

Status as species: Mayr, 1870b: 958 (in key); Forel, 1893g: 351; Dalla Torre, 1893: 169; von Jhering, 1894: 378; Forel, 1895b: 110; Forel, 1899c: 102; Forel, 1905b: 159; Wheeler, W.M. 1905b: 132; Forel, 1907e: 7; Forel, 1908c: 395; Wheeler, W.M. 1911a: 29; Wheeler, W.M. 1911b: 170; Forel, 1912h: 47; Emery, 1913a: 27; Wheeler, W.M. 1913d: 242; Wheeler, W.M. 1917g: 462; Luederwaldt, 1918: 47; Borgmeier, 1923: 87; Störcke, 1926: 117 (in key); Eidmann, 1936b: 89; Wheeler, W.M. 1942: 213; Creighton, 1950a: 342; Kempf, 1972a: 125; Shattuck, 1994: 124; Bolton, 1995b: 247; Wild, 2007a: 67 (redescription); Wild, 2007b: 25; Branstetter & Sáenz, 2012: 254; Bezděčková, *et al.* 2015: 109; Escárraga & Guerrero, 2016: 453; Guerrero, 2019: 714.

Senior synonym of *bicolor*: Wild, 2007a: 67.

Senior synonym of *dominicensis*: Wild, 2007a: 67.

Senior synonym of *fuscescens*: Wild, 2007a: 67.

Senior synonym of *melleus*: Wild, 2007a: 67.

Senior synonym of *nigellus*: Wild, 2007a: 67.

Senior synonym of *succineus*: Wild, 2007a: 67.

Material of the unavailable name *paranaensis* referred here by Wild, 2007a: 67.

keiteli. *Iridomyrmex keiteli* Forel, 1907e: 8 (w.q.) HAITI.

Wild, 2007a: 78 (m.).

Combination in *Linepithema*: Shattuck, 1992a: 16.

Status as species: Emery, 1913a: 27; Wheeler, W.M. & Mann, 1914: 42; Kempf, 1972a: 125; Shattuck, 1994: 125; Bolton, 1995b: 247; Wild, 2007a: 76 (redescription); Lubertazzi, 2019: 126.

Senior synonym of *subfasciatus*: Wild, 2007a: 77.

- leucomelas.** *Iridomyrmex leucomelas* Emery, in von Jhering, 1894: 378 (footnote) (w.q.) BRAZIL (Rio Grande do Sul, Santa Catarina, Rio de Janeiro).
Wild, 2007a: 83 (m.).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Forel, 1908c: 394; Emery, 1913a: 27; Luederwaldt, 1918: 47; Borgmeier, 1923: 88; Kempf, 1969: 294 (redescription); Kempf, 1972a: 126; Shattuck, 1994: 126; Bolton, 1995b: 247; Wild, 2007a: 81 (redescription).
Senior synonym of aspidocoptus: Wild, 2007a: 81.
- melleus.** *Iridomyrmex melleus* Wheeler, W.M. 1908a: 151, pl. 12, figs. 15, 16 (w.q.m.) PUERTO RICO.
Wheeler, G.C. & Wheeler, J. 1974b: 397 (l.).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Emery, 1913a: 27; Smith, M.R. 1937: 863; Kempf, 1972a: 126; Shattuck, 1994: 126; Bolton, 1995b: 247.
Junior synonym of *iniquum*: Wild, 2007a: 68.
- micans.** *Iridomyrmex dispertitus* subsp. *micans* Forel, 1908c: 394 (w.m.) BRAZIL (São Paulo).
Forel, 1911c: 306 (q.).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of dispertitum: Forel, 1911c: 306; Emery, 1913a: 26; Bruch, 1914: 226; Gallardo, 1916a: 106; Wheeler, W.M. 1916d: 330; Luederwaldt, 1918: 47; Wheeler, W.M. 1922c: 14; Borgmeier, 1923: 87; Menozzi, 1931b: 274; Kempf, 1972a: 124; Shattuck, 1994: 122; Bolton, 1995b: 247.
Status as species: Wild, 2007a: 85 (redescription); Wild, 2007b: 25.
Senior synonym of platensis: Wild, 2007a: 85.
Senior synonym of scotti: Wild, 2007a: 85.
Material of the unavailable names *pertaesta*, *transiens* referred here by Wild, 2007a: 85.
- neotropicum.** *Linepithema neotropicum* Wild, 2007a: 91, figs. 6, 29, 30, 65, 104 (w.q.m.) PARAGUAY, BOLIVIA, BRAZIL (Amazonas, Bahia, Espírito Santo, Goiás, Maranhão, Mato Grosso, Minas Gerais, Paraná, Pernambuco, Rio de Janeiro, Rondônia, São Paulo, Tocantins), COLOMBIA, COSTA RICA, GUYANA, TRINIDAD, VENEZUELA.
Status as species: Wild, 2007b: 25; Bezděčková, *et al.* 2015: 109; Escárraga & Guerrero, 2016: 454; Guerrero, 2019: 714.
- nigellus.** *Iridomyrmex iniquus* var. *nigellus* Emery, 1890b: 56 (w.) COSTA RICA.
[Also described as new by Emery, 1894k: 61.]
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *dispertitum*: Forel, 1901c: 134; Forel, 1908b: 62; Forel, 1912h: 47,
Subspecies of *iniquum*: Dalla Torre, 1893: 169; Forel, 1899c: 102; Emery, 1913a: 27; Wheeler, W.M. 1917g: 462; Wheeler, W.M. 1918b: 26; Wheeler, W.M. 1942: 213; Creighton, 1950a: 342; Smith, M.R. 1951a: 836; Smith, M.R. 1958c: 140; Kempf, 1972a: 125; Smith, D.R. 1979: 1419; Shattuck, 1994: 125; Bolton, 1995b: 247; Coover, 2005: 107.
Junior synonym of *iniquum*: Wild, 2007a: 67.
- oblongum.** *Iridomyrmex humilis* var. *oblongus* Santschi, 1929d: 306 (w.) ARGENTINA (Jujuy).
[Misspelled as *oblungus* by Menozzi, 1935c: 321, Goetsch & Menozzi, 1935: 97.]
Wild, 2007a: 98 (q.m.).
Combination in *Linepithema*: Shattuck, 1992a: 16.

- Subspecies of *humile*:** Santschi, 1929d: 307 (in key); Goetsch & Menozzi, 1935: 97; Menozzi, 1935c: 321; Kempf, 1970a: 26; Kempf, 1972a: 125.
- Status as species:** Snelling, R.R. & Hunt, 1976: 90; Shattuck, 1994: 126; Bolton, 1995b: 247; Wild, 2007a: 97 (redescription).
- paranaensis*. *Iridomyrmex melleus* st. *succineus* var. *paranaensis* Santschi, 1929d: 305 (w.) BRAZIL (Paraná); unavailable (infrasubspecific) name.
As unavailable (infrasubspecific) name: Kempf, 1972a: 125.
Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 127.
Material referred to *iniquum* by Wild, 2007a: 68.
- pertaesta*. *Iridomyrmex humilis* st. *angulata* var. *pertaesta* Santschi, 1916e: 390 (w.m.) ARGENTINA (Buenos Aires); unavailable (infrasubspecific) name.
Santschi, 1929d: 307 (q.).
As unavailable (infrasubspecific) name: Santschi, 1929d: 306; Kempf, 1972a: 125.
Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 127.
Material referred to *micans* by Wild, 2007a: 86.
- piliferum*. *Hypoclinea pilifera* Mayr, 1870a: 393 (w.q.) COLOMBIA (“New Granada”).
Crozier, 1970: 113 (k.). Wild, 2007a: 103 (m.).
Combination in *Iridomyrmex*: Dalla Torre, 1893: 169;
combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Mayr, 1870b: 959 (in key); Dalla Torre, 1893: 169; Emery, 1913a: 27; Kempf, 1972a: 126; Shattuck, 1994: 127; Bolton, 1995b: 247; Wild, 2007a: 101 (redescription); Guénard & Economo, 2015: 226; Escárraga & Guerrero, 2016: 455; Guerrero, 2019: 714.
- platensis*. *Iridomyrmex humilis* r. *platensis* Forel, 1912h: 46 (w.q.m.) ARGENTINA (Buenos Aires).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *humile*: Forel, 1913l: 242; Forel, 1914d: 287; Bruch, 1914: 226; Forel, 1915c: 359; Gallardo, 1916a: 101; Santschi, 1916e: 390; Santschi, 1929d: 307 (in key); Kempf, 1972a: 125; Shattuck, 1994: 124; Bolton, 1995b: 247.
Junior synonym of *micans*: Wild, 2007a: 86.
- pordescens*. *Iridomyrmex pordescens* Wheeler, W.M. 1942: 214 (w.) COSTA RICA.
Combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Kempf, 1972a: 126; Shattuck, 1994: 127; Bolton, 1995b: 247.
Junior synonym of *angulatum*: Wild, 2007a: 29.
- praenotandum*. *Iridomyrmex humilis* var. *praenotandum* Schmitz, 1950: 14, PORTUGAL.
Nomen nudum.
- pulex*. *Iridomyrmex humilis* st. *angulata* var. *pulex* Santschi, 1923d: 68 (w.) BRAZIL (Santa Catarina); unavailable (infrasubspecific) name.
As unavailable (infrasubspecific) name: Santschi, 1929d: 308 (in key); Santschi, 1936d: 417; Kempf, 1972a: 125.
Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 127.
- pulex*. *Linepithema pulex* Wild, 2007a: 105, figs. 45, 46, 69, 105 (w.q.m.) PARAGUAY, ARGENTINA (Misiones), BRAZIL (Espírito Santo, Goiás, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina, São Paulo).
Status as species: Wild, 2007b: 25.

- riograndensis*. *Iridomyrmex riograndensis* Borgmeier, 1928b: 64 (w.) BRAZIL (Rio Grande do Sul).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Status as species: Kempf, 1972a: 126; Shattuck, 1994: 127; Bolton, 1995b: 247.
Junior synonym of *humile*: Wild, 2004: 1207; Wild, 2007a: 61; Radchenko, 2016: 84.
- scotti*. *Iridomyrmex humilis* st. *scotti* Santschi, 1919f: 52 (w.) ARGENTINA (Córdoba).
Combination in *Linepithema*: Shattuck, 1992a: 16.
As unavailable (infrasubspecific) name: Santschi, 1929d: 307 (in key).
Subspecies of *humile*: Eidmann, 1936b: 88; Kempf, 1972a: 125; Shattuck, 1994: 124; Bolton, 1995b: 247.
Junior synonym of *micans*: Wild, 2007a: 86.
- subfasciatus*. *Iridomyrmex keiteli* var. *subfasciatus* Wheeler, W.M. & Mann, 1914: 43 (w.) HAITI.
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *keiteli*: Kempf, 1972a: 126; Shattuck, 1994: 125; Bolton, 1995b: 247.
Junior synonym of *keiteli*: Wild, 2007a: 77.
- succineus*. *Iridomyrmex melleus* subsp. *succineus* Forel, 1908c: 396 (w.) BRAZIL (São Paulo).
Combination in *Linepithema*: Shattuck, 1992a: 16.
Subspecies of *melleus*: Emery, 1913a: 27; Santschi, 1929d: 305.
Subspecies of *iniquum*: Forel, 1912h: 47; Luederwaldt, 1918: 47; Borgmeier, 1923: 88; Kempf, 1972a: 125; Shattuck, 1994: 125; Bolton, 1995b: 247.
Junior synonym of *iniquum*: Wild, 2007a: 68.
- transiens*. *Iridomyrmex humilis* r. *platensis* var. *transiens* Forel, 1913l: 242 (w.q.m.) ARGENTINA (Buenos Aires); unavailable (infrasubspecific) name.
As unavailable (infrasubspecific) name: Bruch, 1914: 226; Gallardo, 1916a: 103; Santschi, 1929d: 307 (in key); Kempf, 1972a: 125.
Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 127.
Material referred to *micans* by Wild, 2007a: 86.
- tsachila*. *Linepithema tsachila* Wild, 2007a: 109, figs. 19, 20, 47, 51, 63, 64, 103 (w.q.m.) ECUADOR, COLOMBIA.
Status as species: Escárraga & Guerrero, 2016: 456; Guerrero, 2019: 714.