

Ground foraging ants (Hymenoptera: Formicidae) in Argane (*Argania spinosa*, L) ecosystem: Response to grazing impact

Abdelhadi Ajerrar^{1,2}, Rachid Bouharroud^{1,*}, Mina Zaafrani², Naima Ait Aabd¹, Redouan Qessaoui^{1,2}, Hilal Bahadou², El Hassan Mayad³, Abdelghani Tahiri¹, Bouchra Chebli²

¹Research Unit of Integrated Crop Production, Regional Center of Agronomic Research of Agadir, Morocco

²Biotechnology and Environmental Engineering Unit, National School of Applied Sciences, Ibn Zohr University, Morocco

³Laboratory of Biotechnology and Valorization of Natural Resources Department of Biology, Faculty of Sciences, Ibn Zohr University, BP 8106, 80000 Agadir, Morocco

(Received: January 07, 2020; Revised: April 27, 2020 ; Accepted: June 24, 2020)

ABSTRACT

Grazing is one of the important activities practiced by local farmers in the argane ecosystem. Agdal is the traditional management system linked to grazing management in this ecosystem. This work focused on grazing and Agdal impact on ground foraging ant community. Ant sampling using pitfall traps was occurred in three selected sites during two periods. Species richness, abundance and ant diversity per sample unit were compared between site 1 which is an open rangeland and two sites (2 and 3) where a seasonal defense for grazing and harvesting is applied (Agdal). Ants workers captured were belonging to 14 species, 10 genera and 3 subfamilies. *Monomorium salomonis obscuriceps* (Santschi, 1921) and *Pheidole pallidula* (Nylander, 1849) were the two predominated species during November-January period samples. They contribute to more than 50 % of dissimilarity between ant communities. A negative impact of overgrazing on diversity parameters was reported in site 1. Significant decrease of species richness, abundance and ant diversity per sample unit was observed during both sample periods in overgrazed site (site1) compared to the other sites where Agdal management system is applied.

Key words: Agdal, *Argania spinosa*, Biodiversity, Formicidae, *Pheidole pallidula*, *Temnothorax tameriensis*, *Tapinoma magnum*

INTRODUCTION

Ants are important components of terrestrial ecosystems in terms of biomass and diversity, playing a crucial role in their function (Hölldobler & Wilson, 1990). They live in various environments with diverse feeding habits and in association with other species, in particular, plants and insects (Hölldobler & Wilson, 1990). Ants are important ecologically because of their actions at many trophic levels in the ecosystem - as predators as prey, as detritivores, mutualists, and herbivores (Alonso, 2000). Ant communities appear to be perfect candidates as bio-indicators (Majer, 1983; Nash *et al.*, 2004). According to Andersen & Majer (2004) ants monitoring has been successfully applied to a wide range of other land-use situations. They were used to monitoring ecosystem restoration following mining (Andersen, 1997), forest management (Aman *et al.*, 2009), conservation assessment (Hevia *et al.*, 2013) and grazing impacts (Bestelmeyer & Wiens, 1996; Nash *et al.*, 1999; Nash *et al.*, 2004; Hoffmann, 2010). In Morocco more than 85 % of land is classified as rangeland, which includes forests, steppes, high meadows, and Saharan lands. As a result of intensive grazing without proper management,

the state of Morocco realized that millions of hectares of rangeland are being degraded, especially in the north-eastern area, the *Argania spinosa* ecosystem, and the sub-Saharan and Saharan area (Nash *et al.*, 2014). Souss-valley and its surrounding mountains constitute an exceptional area, where *Argania spinosa* is exclusively endemic (Blérot & Mhirit, 1999). The argane forest has been declared as Biosphere Reserve in 1998. Argane tree has a major role in the ecology, economy, and social relations of the local communities (Ait Aabd *et al.*, 2013; Belyazid, 2000). Local farmers still practice traditional management system in Biosphere Reserve Arganeraie. The agro-sylvo-pastoral system in RBA is based on four major productions: billy goat, barley, wood and argane oil (Bourbouze & El Aïch, 2005). People from the argane region have developed a particular management system to protect fruits during maturation (Bourbouze & El Aïch, 2005). In the argane grove, a seasonal defense for grazing and harvesting is traditionally applied, called Agdal (Bourbouze & El Aïch, 2005). We hypothesise that this traditional management system practice (Agdal) has a positive impact on ants which are an important component of argane ecosystem, they represents the most specious and

*Corresponding Author's E-mail: bouharroud@yahoo.fr




Figure 1. Map of the study sites (Grey spots)

abundant family, it represents more than 50% of total ground foraging arthropods (Ajerrar *et al.*, 2015; Ajerrar *et al.*, 2017). To confirm this hypothesis we suggest to compare ants species richness, abundance, and Shannon diversity between an overgrazing site and two sites where Agdal management system was applied.

MATERIALS AND METHODS

Study sites

This work has been conducted in three Argane ecosystem sites in Souss valley located in south-west of Morocco. The three study sites belong to the argane forest ecosystem. The first site (site 1) is located in Argana (30.7519869N, -9.1529756W), in the north-west of Taroudannt, this site is an open rangeland without any management system; cattle's frequent daily this area. The second site (Site 2) is located in Aoulouz (30.62352N, -8.14542W) in Taroudannt. The third site (Site 3) is located in Tameri (30.70672N, -9.76643W) in the north of Agadir (Figure 1, Table 1). Site 2 and Site 3 are managed plots for barley crops in the shade of argane trees and for argane fruits production. Barley cropping usually starts on first autumnal rain, and harvesting during late spring. The Agdal management system is applied in these two sites; farmers cannot harvest the argane fruits and bring their cattle until after a specific date which everyone knows.

Sampling

Ant sampling was carried out using pitfall traps, the predominant method for studying ant communities in

rangelands and arid area (Agosti & Alonso, 2000; Read & Andersen, 2000; Nash *et al.*, 2004; Hoffmann, 2010). Pitfall traps measures are 16 cm in height and 9 cm in diameter (Gonçalves & Pereira, 2012; Assis *et al.*, 2018). In each site, twelve argane trees were selected separated from each other by 10 to 15 m. At each tree four pitfall traps were installed on four cardinal directions at about 80 cm from the trunk. Ant sampling was occurred during January 2016 to November 2016. Traps were carefully installed with minimal soil and litter disturbance. The traps were filled to 1/3th with salted water (10g/L) mixed with drops of natural soap. The traps were removed to laboratory 24 hours after installation. In the laboratory, ants were separated from other soil fauna and rinsed with tap water to remove plant debris and the rest of the soil; then conserved in ethanol 70%. Ant identifications was made to species and genus level using stereomicroscope and appropriate keys and catalogs (Bolton, 1994; Cagniant, 1996a; Cagniant, 1996b; Cagniant, 1997; Cagniant & Espadaler, 1997; Cagniant & Espadaler, 1998; Cagniant, 2006; Cagniant, 2009; Borowiec, 2014) in the laboratory of entomology, integrated crop production unit, at National Institute of Agronomic Research (INRA). Doubtful specimens were identified by Dr. Henri Cagniant (ex-professor of University Paul Sabatier Toulouse). Voucher specimens are available at INRA Entomology laboratory (Agadir, Morocco).

Data analysis

Due to ant seasonality activity in argane ecosystem proved by El keroumi *et al.* (2012) samples were pooled into two periods; (November-January) and (April-

Table 1. Study sites characteristics

Sites	Locality	Altitude (m)	Climate	Types of Soil	Average Trees density (Ha)	Average tree height (m)	Main plants	Associate plants	Anthropogenic activity / management system
Site 1	Argana	775	Arid	Laomy with weak litter layer	50	5	<i>Rhus pentaphylla</i> , <i>Ziziphus lotus</i> , <i>Launaea arborescens</i>		open rangeland without proper management
Site 2	Aoulouz	784	Arid	Laomy with weak litter layer	80	7	<i>Ziziphus lotus</i> , <i>Launaea arborescens</i>		Barley crops, Grazing/ Agdal
Site 3	Tameri	86	Arid	laomy with relative dense litter layer	100	5	<i>Olea europaea</i> , <i>Genista sp</i> , <i>Drimia sp</i> , <i>Euphorbia officinarum</i> , <i>Ononis natrix</i> .		Barley crops, Grazing/ Agdal

August). Software package Paste 3 (Hammer *et al.*, 2001) was used to compute ant richness (S), ant species abundance (N), and Shannon diversity index (H). To avoid overvalue abundance of species forming trails, abundance classes was assigned according to Read & Andersen (2000); 1=1 ant, 2=2-5 ants, 3=6-20 ants, 4=21-50 ants, 5=51-200 ants and 6>200 ants. Ant data recorded were analyzed by ANOSIM and SIMPER (Bray-curtis index, 9999 permutation) test. The main objectives of this analysis were to assess dissimilarity on ant community between the three sites among the two sampling periods and species percentage contribution on dissimilarity. The number of total expected ant species in each study site among the two periods was computed by Estimate S (Colwell, 2006) based on the number of rare species and 100 time randomization. The mean and SD of species ant richness were computed for the 12 samples units in each site. The expected richness function Mao Tau and 95% CI curve was estimated from 100 randomizations. The obtained data of ant diversity (abundance, richness, Shannon index) were subjected to ANOVA one way at $p < 0.05$. Newman-Keuls post hoc test at 95% confidence limit was used to compare mean value. To avoid impact of seasonality, data obtained from each study site was compared during each period.

RESULTS

Ant fauna

Sampling effort yielded a total of 12.664 ant workers in the three sites during the two samples periods. Ants workers captured are belonging to 14 species, 10 genera and 3 subfamilies. Myrmicinae (9 species) and Formicidae (4 species) were the most speciose subfamily whereas Dolichoderinae represented only by one species. Up to 67 % of total ants were captured during April-August period however less than 33% were captured during November-January period. *Monomorium salomonis obscuriceps* was the most abundant species in both sites 1 and 3 however *Pheidole pallidula* was the most

abundant species in site 2. Ant species per sample unit (tree) varied from 0 species in site 1 for only one sample unit to 9 species in site 3. The expected ant richness curves obtained from 100 randomization of samples order tend to decrease with adding samples, therefore sampling effort was sufficient (expected ant richness were similar to observed richness in almost all samples in the three studied sites) (Figure 2). Ant communities of the three studied sites were different with some overlap during November-January period, where ant community of site 2 and 3 shows a low dissimilarity ($R=0.28$; $P=0.0002$), however a relatively high dissimilarity of ant community was reported between site 1 and 3 ($R=0.44$; $P=0.0001$). During April-August period, ant community of the three studied sites was different to highly different without overlap, thus, the highest dissimilarity were reported between site 2 and site 3 ($R=0.96$; $P=0.0001$), as well as between site 2 and site 1 ($R=0.7$; $P=0.0001$) (Table 2). *Monomorium salomonis obscuriceps* and *Pheidole pallidula* contribute to more than 50 % of dissimilarity between ant communities during November-January period. However, a similar dissimilarity percentage contribution of several ant species was shown during April-August period, nevertheless, the high contributed dissimilarity percentage was computed for *Aphaenogaster praedo* Emery, 1908 (15.20 %) and for *Pheidole pallidula* (14.36 %) (Table 3).

ANOVA one way ($p=0.05$) analysis of mean species richness, Shannon diversity and mean species abundance computed per sample unit (tree) showed a significant difference comparing mean computed in each site and during the tow sampling periods. Mean species richness (November-January: $F=16.95$, df. 33, $P<0.00001$; April-August: $F=16.53$, df. 33, $P<0.00001$). Low species richness was recorded in overgrazed site (site1) whatever the sampling periods compared to managed sites (site 2 and 3). Therefore, Newman-Keuls post hoc test (at 95% limit of confidence) showed a significant difference between site 1 and the two other sites (Figure 3). Low abundance was recorded in site 1 during both periods;




Figure 2. Expected ant richness (mean \pm SD) function Mao Tau computed among two periods at each study site.

November-January (2.41 ± 1.6) and April-August (9.42 ± 4), however, a higher abundance was recorded in site 3 during April-August (23.58 ± 4.6). A high and similar abundance value was recorded in both sites 2 and 3 during Autumn-January. Significant difference among the three sites was reported comparing mean abundance (Autumn-January: $F = 28.32$, df. 33, $P < 0.000001$; April-August: $F = 41.55$, df. 33, $P < 0.0001$) (Figure 3). Low ant diversity was reported during November-January period in three studied sites; however relative increase was shown during April-August period. Significant difference between site 1 and the other sites were reported comparing Shannon diversity index (Autumn-January: $F = 21.82$, df. 33, $P < 0.00001$; April-August: $F = 17.51$, df. 33, $P < 0.00001$). The lowest value of Shannon diversity

index was recorded in site 1 (0.39 ± 0.3) during November-January, while a significant increasing was observed during April-August (1.22 ± 0.40) in the same site (Figure 3).

DISCUSSION

Ant ground foraging below argane trees in Souss valley were composed of 14 species. *Temnothorax tamriensis* was the new ant species reported in Morocco (Ajerrar *et al.*, 2018). All captured species are native and no invasive species has been captured. Low ant richness reported in this study was reported elsewhere (Cerdá *et al.*, 2009; El Keroumi *et al.*, 2012; Gonçalves & Pereira, 2012; Barech *et al.*, 2018). Diversity parameters computed in

Table 2. Similarity computed among seasons of ant communities between study sites. ANOSIM R values and their corresponding P (between brackets).

Sites	November-January		April -August	
	site 2	site 3	site 2	site 3
Site 1	0.3496 (0.0001)	0.4469 (0.0001)	0.5184 (0.0001)	0.7 (0.0001)
Site 2		0.2822 (0.0002)		0.9672 (0.0001)

Table 3. average dissimilarity and contribution percentage computed by SIMPER of ant species among study sites during each period.

Taxon	November-January		April-August	
	Av. dissim	Contrib. %	Av. dissim	Contrib. %
<i>Monomorium salomonis obscuriceps</i>	20.28	29.59	6.18	10.54
<i>Pheidole pallidula</i>	16.07	23.44	8.43	14.36
<i>Aphaenogaster praedo</i>	12.29	17.93	8.92	15.20
<i>Messor maroccanus</i>	6.27	9.14	5.02	8.56
<i>Tapinoma magnum</i>	4.01	5.85	6.33	10.79
<i>Tetramorium semilaeve depressum</i>	2.97	4.33	3.98	6.78
<i>Camponotus brullei</i>	1.89	2.76	3.73	6.35
<i>Cataglyphis albicans vaucheri</i>	1.83	2.66	5.39	9.18
<i>Camponotus erigenes</i>	1.50	2.19	6.03	10.28
<i>Temnothorax sp1</i>	0.83	1.21	0.38	0.64
<i>Temnothorax tameriensis</i>	0.61	0.90	2.08	3.54
<i>Temnothorax sp2</i>	0.00	0.00	0.33	0.57
<i>Cataglyphis viatica</i>	0.00	0.00	0.38	0.65
<i>Crematogaster scutellaris algerica</i>	0.00	0.00	1.51	2.57

(Av. dissim= average dissimilarity; Contrib.%= % of contribution)

each site were different between the two sampling periods. Indeed, a high value was computed during warm and dry season (April-August), however, lower values were computed during wet and cold season (November-April). The same finding was reported by El Keroumi *et al.* (2012) study and within the same site the difference may be explained by the seasonality pattern of ground ant foraging activity under argane tree. The obtained results show a significant impact of overgrazing on ant diversity, species richness and abundance per sample unit. Similar studies conducted in arid climate elsewhere reported the negative impact of grazing on ant richness (Nash *et al.*, 2001), richness and relative abundance of some species (Bestelmeyer & Wiens, 2001), abundance (Debinski *et al.*, 2011) and species diversity (Abensperg-Traun *et al.*, 1996). Moreover, grazing impact on ant is worldwide proved in different ecosystems (Hoffmann, 2010). The site 1 located in Argana arganeraie which is overgrazed without any management system shows a significant decrease in abundance, richness and diversity per sample unit compared to the other two sites where traditional management is applied. This significant decrease of the three studied diversity parameters recorded in the site 1 could be explained by alteration of the vegetal cover due to intensive grazing. According to Andersen (1995), grazing disturbance acts indirectly through change in vegetation structure, food supplies and

competitive interactions. In addition, the ripe and ripening argane fruits are known as suitable environment for insect's development and are heavily infected mainly by Mediterranean fruit fly *Ceratitis capitata* (Debouzie & Mazih, 1999). The fact that these fruits are consumed by livestock for its pulp (Bourbouze & El Aïch, 2005) before emergence of insect's larva may reduce ant's food resource availability and impacts negatively their population size and diversity. According to El Keroumi *et al.* (2010), 96% of *C. capitata* larva were captured by ants under argane trees in Essouira argane forest.

Based on different ant studies conducted in Mediterranean ecosystems and our field observations, the ant species associated to arganeraie ecosystem can be classified into three main groups according to feeding behavior: generalist/omnivorous species, seed feeder, and insect, honeydew and nectar feeder. Therefore, four species (*P. pallidula*, *T. semilaeve*, *M. salomonis*, *A. praedo*) are belonging to generalist feeders. However, *M. maroccanus* is only a seed / generalist feeder. Three species are belonging to insectivore/animal feeder (*C. albicans*, *C. viatica* and *C. scutellaris*). In addition, *T. magnum* was shown to visit Argane trees for Aphid's honeydew and argane flower nectar (Ajerrar *et al.*, 2020). *Pheidole P. pallidula* and *T. semilaeve* are two omnivorous ant that collect animal and plant remains in similar proportion and that rarely climb the plants to collect liquid food




Figure.3. Ant species richness, ant abundance and Shannon's diversity index (mean values \pm SD) computed during the two sampling periods at each study site. The letters on the graphs denote the significant differences (Newman-keuls post hoc test at 95% limit confidence).

(Retana *et al.*, 1992). *M. salomonis* and *A. praedo* were observed to prey in *C. capitata* larva and plant (per obs). In contrast, Segev & Ziv (2012) study state that *M. salomonis* is a generalist seed-feeder. According to Santini *et al.* (2011) *C. scutellaris* is a common ant widely distributed in both natural and managed ecosystems throughout the Mediterranean area, it occupied a high rank of dominance that has a high ability to discover and to monopolize food sources (Santini *et al.*, 2007).

CONCLUSION

Despite the resilience of ant communities in arid area to disturbance noted elsewhere (Nash *et al.*, 1998; Whitford *et al.*, 1999) overgrazing occurred in open rangeland of argane ecosystem impact negatively richness, abundance and diversity of ants. In the other hand, Agdal management system adopted by local farmers either in their own orchards or in collective ones shows the importance of this traditional management to preserve ant communities in argane ecosystem.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Agronomic Research, Regional Center of Agronomic Research of Agadir, Morocco (Grant id: PRMT Arganier 2017-20). We would like to acknowledge Professor Henri Cagniant (Toulouse, France) for ant species identification.

REFERENCES

- Abensperg-Traun, M., Smith, G.T., Arnold, G.W. and Steven, D.E. 1996. The effects of habitat fragmentation and livestock-grazing on animal communities in a remnant of gimlet Eucalyptus salubris woodland in the Western Australian wheat belt. I. Arthropods. *Journal of Applied Ecology*, 33:1281–1301.
- Agosti, D. and Alonso, L.E. 2000. The ALL Protocol: standard protocol for the collection of ground-dwelling ants. Pp. 204–206. In *Ants: standard methods for measuring and monitoring biodiversity* (eds Agosti, D., Majer, J.D., Alonso, L.E. and Schultz, T.R.), DC: Smithsonian Institution Press, Washington.
- Ait Aabd, N., Bouharroud, R., Tahiri, A., Wifaya, A., Mimouni, A. and El Mousadik, A. 2019. Genetic Diversity, Conservation and Breeding of Argan Tree (*Argania spinosa* L. Skeels). Pp. 31–56. In *Advances in Plant Breeding Strategies: Nut and Beverage Crops* (eds Al-Khayri, J.M., Jain, S.M. and Johnson, D.V.), Springer Nature, Switzerland.
- Ajerrar, A., Bouharroud, R., Amarrague, A., Qessaoui, R. and Mayad, E. 2015. Evaluation de la biodiversité de l'entomofaune terrestre associée à l'écosystème de l'arganier. In: *Proceedings of 3ème congrès international de l'arganier* (vol. 3, pp. 102-105), Agadir, 17-19 December 2015.

- Ajerrar, A., Bouharroud, R., Zaafrani, M., Chebli, B., Amarraque, A., Qessaoui, R. and Mayad, E. 2017. Abondance de la classe des insectes associés à l'écosystème de l'arganier du Souss. In: Proceedings of 4ème congrès international de l'arganier (vol. 4, pp. 239-244), Agadir, 20-22 November 2017.
- Ajerrar, A., Gomez, K., Bouharroud, R., Zaafrani, M. and Cagniant, H. 2018. *Temnothorax tamriensis* a new species of ant from Morocco. *Revue de l'Association Roussillonnaise d'Entomologie*, Tome XXVII (2): 97-107.
- Ajerrar, A., Akroud, H., Ait Aabd, N., Qessaoui, R., Amarraque, A., Lahmyd, H., Zaafrani, M., Chebli, B., Mayad, El H. and Bouharroud, R. 2020. Pollination system and effective pollinators of *Argania spinosa* (L. Skeels), Journal of the Saudi Society of Agricultural Sciences doi.org/10.1016/j.jssas.2020.04.002.
- Alonso, L.E. 2000. Ants as indicators of diversity. Pp. 80-88. In *Ants: standard methods for measuring and monitoring biodiversity* (eds Agosti, D., Majer, J.D., Alonso, L.E. and Schultz, T.R.), DC: Smithsonian Institution Press, Washington.
- Andersen, A.N. 1995. A classification of Australian ant communities based on functional groups which parallel plant life-forms in relation to stress and disturbance. *Journal of Bio-geography*, 22: 15-29.
- Andersen, A.N. 1997. Ants as indicators of ecosystem restoration following mining a functional group approach. Pp. 319-25. In *conservation outside nature reserves*. (eds Hale, P. and Lamb, D.), Centre For Conservation Biology, The University of Queensland, Brisbane.
- Andersen, A.N., Fisher, A., Hoffmann, B.D., Read, J.L. and Richards, R. 2004. The use of terrestrial invertebrates for biodiversity monitoring in Australian rangelands, with particular reference to ants. *Austral Ecology*, 29: 87-92.
- Andersen, A.N. and Majer, J.D. 2004. Ants show the way Down Under: invertebrates as bioindicators in land management. *Frontiers in Ecology and the Environment*, 2: 291-298.
- Aman, X., Gracia, M., Comas, L. and Retana, J. 2009. Forest management conditioning ground ant community structure and composition in temperate conifer forests in the Pyrenees Mountains. *Forest Ecology and Management*, 258: 51-59.
- Assis, D.S., Dos Santos, I.A., Ramos, F.N., Barrios-Rojas, K.E., Majer, J.D. and Vilela, E.F. 2018. Agricultural matrices affect ground ant assemblage composition inside forest fragments. *PLoS ONE*, 13: e0197697.
- Barech, G., Khaldi, M., Boujelal, F.Z. and Espadaler, X. 2018. Diversité et structure de la myrmécofaune aux abords du barrage El ksob en Algérie: nouvelle citation pour *Aphaenogaster rupestris* forel, 1909 (Hymenoptera: Formicidae). *Boletín de la Sociedad Entomológica Aragonesa*, 62: 253-258.
- Belyazid S (2000). Achieving sustainability in the Argan Forest, Morocco (Unpublished master's thesis). Lund University, Sweden.
- Bestelmeyer, B.T. and Wiens, J.A. 1996. The effects of land use on the structure of ground foraging ant in the Argentine Chaco. *Ecological Applications*, 6: 1225-1240.
- Bolton, B. (1994). Identification guide to the ant genera of the World. Harvard University Press. Cambridge.
- Bestelmeyer, B.T. and Wiens, J.A. 2001. Ant biodiversity in semiarid landscape mosaics: the consequences of grazing vs. natural heterogeneity. *Ecological Applications*, 11: 1123-1140.
- Borowiec, L. 2014. Catalogue of ants of Europe, the Mediterranean Basin and adjacent regions (Hymenoptera: Formicidae). *Biologica Silesiae*. wrocław.
- Bourbouze, A. and El Aïch, A. 2005. L'élevage caprin dans l'arganeraie: l'utilisation conflictuelle d'un espace. *Cahiers Agricultures*, 14: 447-453.
- Cagniant, H. 1996a. Les *Aphaenogaster* du Maroc (Hymenoptera: Formicidae): clé et catalogue des espèces. *Annales de la Société Entomologique de France*, 32: 67-85.
- Cagniant, H. 1996b. Les *Camponotus* du Maroc (Hymenoptera: Formicidae): clé et catalogue des espèces. *Annales de la Société Entomologique de France*, 32: 87-100.
- Cagniant, H. 1997. Le genre *Tetramorium* au Maroc (Hymenoptera: Formicidae): clé et catalogue des espèces. *Annales de la Société Entomologique de France*, 33: 89-100.
- Cagniant, H. 2005. Les *Crematogaster* du Maroc (Hym., Formicidae). Clé de détermination et commentaires. *Orsis*, 20: 7-12.
- Cagniant, H. 2006. Liste actualisée des fourmis du Maroc (Hymenoptera: Formicidae). *Myrmecologische Nachrichten*, 8: 193-200.
- Cagniant, H. 2009. Le Genre *Cataglyphis* Förster, 1850 au Maroc (Hyménoptères: Formicidae). *Orsis*, 24: 41-71.
- Cagniant, H. and Espadaler, X. 1997a. Le genre *Messor* au Maroc (Hymenoptera: Formicidae). *Annales de la Société Entomologique de France*, 33: 419-434.
- Cagniant, H. and Espadaler, X. 1997b. Les *Leptothorax*, *Epimyrma* et *Chalepoxenus* du Maroc (Hymenoptera: Formicidae). Clé et catalogue des espèces. *Annales de la Société Entomologique de France*, 33: 259-284.
- Cerdá, X., Palacios, R. and Retana, J. 2009. Ant Community Structure in Citrus Orchards in the Mediterranean Basin: Impoverishment as a Consequence of Habitat Homogeneity. *Environmental Entomology*, 38: 317-324.
- Collins, S.L. and Smith, M.D. 2006. Scale-dependent interaction of fire and grazing on community heterogeneity in tall grass prairie. *Ecology*, 87: 2058-2067.
- Colwell, R.K. 2006. EstimateS: statistical estimation of species richness and shared species from samples. Version 8.0 Users guide and application. Available online at <http://viceroy.eeb.uconn.edu/Estimates>.

- Debinski, D.D., Moranz, R. A., Delaney, J. T., Miller, J. R., Engle, D. M., Winkler, L. B., McGranahan, D. A., Bamey, R. J., Trager, J. C., Stephenson, A. L. and Gillespie, M. K. 2011. A cross-taxonomic comparison of insect responses to grassland management and land-use legacies. *Ecosphere*, 2:131. doi: 10.1890/ES11-00226.1
- Debouzi, D. and Mazih, A. 1999. Infestation rate of argan fruit (*Argania spinosa*) by the Mediterranean fruit fly (*Ceratitis capitata*) in relation to phenology and maturation of the fruit. *Entomologia Experimentalis et Applicata*, 81: 31-38.
- El Keroumi, A., Naamani, K., Dahbi, A., Luquec, I., Carvajal, A., Cerda, X. and Boulay, R. 2010. Effect of ant predation and abiotic factors on the mortality of medfly larvae, *Ceratitis capitata*, in the Argan forest of Western Morocco. *Biocontrol Science and Technology*, 20: 751-762
- El Keroumi, A., Naamani, K., Soummane, H. and Dahbi, A. 2012. Seasonal dynamics of ant community structure in the Moroccan Argan Forest. *Journal of Insect Science*, 12:94. Available online: insect-science.org/12.94.
- Folgarait, P.J. 1998. Ant biodiversity and its relationship to ecosystem functioning: a review. *Biodiversity and Conservation*, 7: 1221-1244.
- Gomez, C., Casellas, D., Oliveras, J. and Bas, M.J. 2003. Structure of ground-foraging ant assemblages in relation to land-use change in the northwestern Mediterranean region. *Biodiversity and Conservation*, 12: 2135-2146.
- Gonçalves, M.F. and Pereira, J.A. 2012. Abundance and diversity of soil arthropods in the olive grove ecosystem. *Journal of Insect Science*, 12:20. Available online: insectscience.org/12.20
- Hammer, Ø., Harper, D.A.T. and Ryan, P.D. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, 4: 9pp.
- Hevia, V., Azcarate, F.M., Oteros-Rozas, E. and González, J.A. 2013. Exploring the role of transhumance drove roads on the conservation of ant diversity in Mediterranean agroecosystems. *Biodiversity and Conservation*, 22:2567-2581
- Hoffmann, B.D. 2010. Using ants for rangeland monitoring: Global patterns in the responses of ant communities to grazing. *Ecological Indicators*, 10: 105-111.
- Hölldobler, B. and Wilson, E.O. 1990. The ants. Springer. Berlin.
- Izhaki, I., Levey, D.J. and Silva, W.R. 2003. Effects of prescribed fire on an ant community in sifting and pitfall trapping, southeast. *Naturalist*, 3: 113-126.
- Living, S.C. 1983. Seasonal, annual, and among-site variation in the ground ant community of a deciduous tropical forest: Some causes of patchy species distributions. *Ecological Monographs*, 53: 435-455.
- Majer, J.D. 1983. Ants: Bio-indicators of mine site rehabilitation, Land-Use, and Land Conservation. *Environmental Management*, 7: 375-383.
- Nash, M.S., Whitford, W.G., Van Zee, J. and Havstad, K. 1998. Monitoring changes in stressed ecosystems using spatial patterns of ant communities. *Environmental Monitoring and Assessment*, 51: 201-210.
- Nash, M.S., Whitford, W.G., deSoyza, A.G., Van Zee, J.W. and Havstad, K.M. 1999. Livestock activity and Chihuahuan Desert annual-plant communities: Boundary analysis of disturbance gradients. *Ecological Applications*, 9: 814-823.
- Nash, M.S., Whitford, W.G., Bradford, D.F., Franson, S.E., Neale, A.C. and Heggem, D.T. 2001. Ant communities and livestock grazing in the Great Basin, USA. *Journal of Arid Environments*, 49: 695-710.
- Nash, M.S., Bradford, D.F., Franson, S.E., Neale, A.C., Whitford, W.G. and Heggem, D.T. 2004. Livestock grazing effects on ant communities in the eastern Mojave Desert, USA. *Ecological Indicators*, 4: 199-213.
- Nash, M.S., Chaloud, D.J., Kepner, W.G. and Sarri, S. 2008. Regional Assessment of Landscape and Land Use Change in the Mediterranean Region. Pp.143-165. In Environmental change and human security: recognizing and acting on hazard impacts. (eds Liotta, P.H., Mouat, D.A., Kepner, W.G. and Lancaster, J.M.), Springer Netherlands, Dordrecht.
- Retana, J., Cerdá, X. and Espadaler, X. 1992. Coexistence of two sympatric ant species, *Pheidole pallidula* and *Tetramorium semilaeve* (Hymenoptera: Formicidae). *Entomologia Generalis*, 17: 29 - 40.
- Read, J.L. and Andersen, A.N. 2000. The value of ants as early warning bioindicators: responses to pulsed cattle grazing at an Australian arid zone locality. *Journal of Arid Environments*, 45: 231-251.
- Santini, G., Tucci, L., Ottonetti, L. and Frizzi F. 2007. Competition trade-offs in the organisation of a Mediterranean ant assemblage. *Ecological Entomology*, 32: 319-326.
- Santini, G., Ramsay, P.M., Tucci, L., Ottonetti, L. and Frizzi, F. 2011. Spatial patterns of the ant *Crematogaster scutellaris* in a model ecosystem. *Ecological Entomology*, 36: 625-634.
- Segev, U. and Ziv, Y. 2012. Consequences of behavioral vs. numerical dominance on foraging activity of desert seed-eating ants. *Behavioral Ecology and Sociobiology*, 66: 623-632.
- Whitford, W.G., Van Zee, J., Nash, M.S., Smith, W.E. and Herrick, J.E. 1999. Ants as indicators of exposure to environmental stressors in North American desert grasslands. *Environmental Monitoring and Assessment*, 54: 143-171.

