

FORELIUS

albiventris. *Forelius chalybaeus* r. *albiventris* Forel, 1912h: 45 (w.) ARGENTINA (Catamarca).

Cuezzo, 2000: 216 (q.m.l.).

Subspecies of chalybaeus: Forel, 1913l: 241; Forel, 1914d: 285; Bruch, 1914: 226; Gallardo, 1916a: 89; Bruch, 1916: 344; Gallardo, 1918a: 31.

Status as species: Kusnezov, 1953b: 339; Kusnezov, 1954b: 37; Kusnezov, 1957b: 16 (in key); Kempf, 1972a: 108; Shattuck, 1994: 92; Bolton, 1995b: 190; Cuezzo, 2000: 215 (redescription).

anale. *Tapinoma analis* André, 1893b: 148 (w.) MEXICO (Chihuahua).

Combination in *Iridomyrmex*: Forel, 1908b: 62;

combination in *Forelius*: Wheeler, G.C. & Wheeler, J. 1986g: 13.

Subspecies of pruinosus: Pergande, 1894: 163; Emery, 1895c: 333; Wheeler, W.M. 1905b: 132; Emery, 1913a: 26; Wheeler, W.M. 1913b: 497; Wheeler, W.M. 1913c: 116; Smith, M.R. 1930a: 5; Wheeler, W.M. 1934f: 142; Wheeler, W.M. 1935g: 36; Dennis, 1938: 294; Wesson, L.G. & Wesson, R.G. 1940: 100; Cole, 1942: 373; Buren, 1944a: 291; Creighton, 1950a: 343; Smith, M.R. 1958c: 140; Cole, 1966: 18; Beck, et al. 1967: 70; Smith, M.R. 1967: 364; Kempf, 1972a: 126; Alayo, 1974: 42; Hunt & Snelling, 1975: 22; Yensen, et al. 1977: 183; Smith, D.R. 1979: 1419; Shattuck, 1994: 96.

Status as species: Forel, 1899c: 101; Forel, 1908b: 62; Wheeler, W.M. 1906d: 342; Forel, 1909a: 250; Wheeler, W.M. 1910g: 569; Santschi, 1911d: 7; Emery, 1913a: 26; Cole, 1937b: 136; Smith, M.R. 1951a: 835; Mackay, Lowrie, et al. 1988: 102; Cuezzo, 2000: 222 (redescription); Mackay & Mackay, 2002: 257.

Junior synonym of pruinosus: Snelling, R.R. & George, 1979: 166; Wheeler, G.C. & Wheeler, J. 1986g: 13; Bolton, 1995b: 190; Covert, 2005: 108; Ward, 2005: 9.

andinus. *Forelius andinus* Kusnezov, 1957b: 15 (w.q.) ARGENTINA (Salta).

Cuezzo, 2000: 226 (m.).

Status as species: Kempf, 1972a: 108; Shattuck, 1994: 92; Bolton, 1995b: 190; Cuezzo, 2000: 224 (redescription).

andrei. *Forelius maccooki* r. *andrei* Forel, 1912h: 44 (w.) U.S.A. (California).

Subspecies of foetida: Smith, M.R. 1951a: 836.

Junior synonym of foetida: Wheeler, W.M. 1901b: 520; Snelling, R.R. & George, 1979: 164.

Junior synonym of mccooki: Smith, D.R. 1979: 1419; Bolton, 1995b: 190.

bahianus. *Forelius bahianus* Cuezzo, 2000: 227, figs. 12, 35 (w.q.) BRAZIL (Bahia).

basalis. *Forelius chalybaeus* st. *grandis* var. *basalis* Santschi, 1922b: 375 (w.q.m.) ARGENTINA (Catamarca); unavailable (infrasubspecific) name.

Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 97.

[*Forelius grandis* var. *basalis* Kempf, 1972a: 109. Unavailable name (variety proposed after 1960).]

Material referred to *grandis* by Cuezzo, 2000: 242.

brasiliensis. *Iridomyrmex (Forelius) maccooki* var. *brasiliensis* Forel, 1908c: 396 (w.) BRAZIL (Rio Grande do Sul).

[Misspelled as *brasiliiana* by Bruch, 1914: 226.]

Wheeler, G.C. & Wheeler, J. 1966: 730 (l.). Cuezzo, 2000: 231 (q.m.).

Combination in *Forelius*: Emery, 1913a: 36.

Subspecies of *mccooki*: Forel, 1909a: 260; Forel, 1912h: 43; Forel, 1913l: 241; Bruch, 1914: 226; Gallardo, 1916a: 78; Santschi, 1916e: 391; Wheeler, W.M. 1942: 215.

Subspecies of *foetida*: Emery, 1913a: 36.

Status as species: Santschi, 1922b: 373; Borgmeier, 1923: 98; Kusnezov, 1957b: 17 (in key); Kempf, 1972a: 108; Zolessi, *et al.* 1988: 7; Shattuck, 1994: 92; Bolton, 1995b: 190; Cuezzo, 2000: 229 (redescription); Wild, 2007b: 24.

Senior synonym of *fiebrigi*: Cuezzo, 2000: 230.

Senior synonym of *pilipes*: Cuezzo, 2000: 229.

Material of the unavailable name *carmelitana* referred here by Cuezzo, 2000: 229.

***breviscapus*.** *Forelius maccoki* r. *breviscapus* Forel, 1914d: 286.

[First available use of *Forelius maccoki* r. *fiebrigi* var. *breviscapa* Forel, 1913l: 241 (w.) ARGENTINA (Buenos Aires, Santa Fe); unavailable (infrasubspecific) name.]
Cuezzo, 2000: 237 (q.m.).

As unavailable (infrasubspecific) name: Bruch, 1914: 226.

Subspecies of *maccoki*: Forel, 1915c: 359; Bruch, 1915: 534; Gallardo, 1916a: 80.

Status as species: Santschi, 1922b: 374; Kusnezov, 1953b: 339; Kusnezov, 1957b: 16 (in key); Kempf, 1972a: 109; Shattuck, 1994: 92; Bolton, 1995b: 190; Cuezzo, 2000: 235 (redescription).

Senior synonym of *obscurata*: Cuezzo, 2000: 235.

Senior synonym of *instabilis*: Cuezzo, 2000: 235.

***carmelitana*.** *Forelius maccoki* subsp. *brasiliensis* var. *carmelitana* Gallardo, 1916a: 79, fig. 35 (w.) URUGUAY; unavailable (infrasubspecific) name.

Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 97.

[*Forelius brasiliensis* var. *carmelitanus* Kempf, 1972a: 108. Unavailable name (variety proposed after 1960).]

Material referred to *brasiliensis* by Cuezzo, 2000: 229.

***chalybaeus*.** *Forelius chalybaeus* Emery, 1906c: 176 (w.q.m.) ARGENTINA (Chubut, Mendoza).

Status as species: Forel, 1912h: 45; Emery, 1913a: 35; Forel, 1913l: 241; Forel, 1914d: 285; Bruch, 1914: 226; Gallardo, 1916a: 84; Bruch, 1916: 341; Gallardo, 1918a: 30; Gallardo, 1919b: 248; Santschi, 1922b: 375; Kusnezov, 1953a: 110 (in key); Kusnezov, 1953b: 339; Kusnezov, 1957b: 17 (in key); Kusnezov, 1960b: 361; Kempf, 1972a: 109; Shattuck, 1994: 93; Bolton, 1995b: 190; Cuezzo, 2000: 239 (redescription).

Senior synonym of *minor*: Cuezzo, 2000: 239.

Senior synonym of *paucistricta*: Cuezzo, 2000: 239.

Senior synonym of *personata*: Cuezzo, 2000: 239.

***damiani*.** *Forelius damiani* Guerrero & Fernández, 2008: 53, figs. 1-3 (w.) COLOMBIA, COSTA RICA, U.S.A. (Texas).

Status as species: Branstetter & Sáenz, 2012: 254; Guerrero, 2019: 710.

***debenedettii*.** *Forelius chalybaeus* st. *debenedettii* Santschi, 1919f: 53 (w.) ARGENTINA (Jujuy).

Subspecies of *grandis*: Kusnezov, 1957b: 18; Kempf, 1972a: 109; Shattuck, 1994: 94; Bolton, 1995b: 190.

Junior synonym of *grandis*: Cuezzo, 2000: 242.

eidmanni Goetsch, 1933; see under **TAPINOMA**.

***fiebrigi*.** *Forelius maccoki* r. *fiebrigi* Forel, 1912h: 44 (w.) PARAGUAY.

Subspecies of *maccooki*: Forel, 1913l: 241; Bruch, 1914: 226; Bruch, 1915: 534; Gallardo, 1916a: 80; Wheeler, W.M. 1942: 215.

Status as species: Kempf, 1972a: 109; Shattuck, 1994: 93; Bolton, 1995b: 190.

Junior synonym of *brasiliensis*: Cuezzo, 2000: 230.

foetida. *Formica foetida* Buckley, 1866: 167 (w.q.) U.S.A. (Texas).

[Junior primary homonym of *Formica foetida* Linnaeus, 1758: 582 (Emery, 1921a: 26).]

Crozier, 1970: 113 (k.); Wheeler, G.C. & Wheeler, J. 1974b: 401 (l.).

Status as species: Cresson, 1887: 256; Dalla Torre, 1893: 196; Wheeler, W.M. 1901b: 520; Emery, 1913a: 35; Creighton, 1950a: 345; Smith, M.R. 1951a: 836; Smith, M.R. 1967: 365; Kempf, 1972a: 109; Hunt & Snelling, 1975: 22; Yensen, *et al.* 1977: 183; Smith, D.R. 1979: 1419; Snelling, R.R. & George, 1979: 163; Allred, 1982: 461; Wheeler, G.C. & Wheeler, J. 1986g: 56.

Senior synonym of *maccooki*: Wheeler, W.M. 1901b: 520; Wheeler, W.M. 1902f: 23; Emery, 1913a: 35; Creighton, 1950a: 345; Smith, M.R. 1951a: 836; Kempf, 1972a: 109; Shattuck, 1994: 94.

Replacement name: *Iridomyrmex mccooki* McCook, 1880: 187.

[Note: *mccooki* McCook junior synonym of *foetida* Buckley (synonymy by Wheeler, W.M. 1901b: 520); hence *mccooki* first available replacement name for *foetida* Buckley.]

golbachi Kusnezov, 1953; see under **LEPTANILLOIDES**.

grandis. *Forelius chalybaeus* r. *grandis* Forel, 1912h: 46 (w.) ARGENTINA (Catamarca).

[Note: Santschi, 1916e: 393, erroneously refers to this taxon as *major*.]

Forel, 1914d: 285 (q.m.).

Subspecies of chalybaeus: Forel, 1913l: 241; Forel, 1914d: 285; Bruch, 1914: 226; Bruch, 1915: 533; Bruch, 1916: 345; Gallardo, 1916a: 87; Santschi, 1916e: 391; Bruch, 1917d: 432; Gallardo, 1918a: 31; Gallardo, 1919b: 249; Santschi, 1922b: 375.

Status as species: Kusnezov, 1954b: 37; Kusnezov, 1957b: 18 (in key); Kempf, 1972a: 109; Shattuck, 1994: 94; Bolton, 1995b: 190; Cuezzo, 2000: 242 (redescription).

Senior synonym of debenedettii: Cuezzo, 2000: 242.

Material of the unavailable names *basalis*, *symbiotica* referred here by Cuezzo, 2000: 242.

instabilis. *Forelius maccooki* st. *instabilis* Santschi, 1920d: 382 (w.q.m.) ARGENTINA (Buenos Aires).

Junior synonym of *obscuratus*: Santschi, 1922b: 374.

Status as species: Kusnezov, 1957b: 16 (in key); Kempf, 1972a: 109; Shattuck, 1994: 94; Bolton, 1995b: 190.

Junior synonym of *breviscapus*: Cuezzo, 2000: 235.

keiferi. *Forelius foetidus* subsp. *keiferi* Wheeler, W.M. 1934f: 141 (w.q.m.) MEXICO (Revillagigedo Is.).

Subspecies of foetida: Wheeler, W.M. 1935g: 36; Kempf, 1972a: 109.

Subspecies of mccooki: Shattuck, 1994: 94; Bolton, 1995b: 190.

Status as species: Cuezzo, 2000: 246 (redescription).

lilloi. *Forelius lilloi* Cuezzo, 2000: 248, figs. 18, 34 (w.) ARGENTINA (Tucumán).

macrops. *Forelius macrops* Kusnezov, 1957b: 14 (w.) ARGENTINA (Tucumán, Salta, Jujuy).

Status as species: Kempf, 1972a: 109; Shattuck, 1994: 95; Bolton, 1995b: 190; Cuezzo, 2000: 253 (redescription).

major, misspelling, see under *grandis*.

maranhaoensis. *Forelius maranhaoensis* Cuezzo, 2000: 255, figs. 11, 27, 38 (w.) BRAZIL (Maranhão, Bahia, Distrito Federal, Goiás, Pernambuco).

mccooki. *Iridomyrmex mccooki* McCook, 1880: 187 (w.) U.S.A. (Texas).

[Misspelled as *maccooki* by Mayr, 1886d: 432, Dalla Torre, 1893: 168, and others.]

Replacement name for *Formica foetida* Buckley, 1866: 167. [Junior primary homonym of *Formica foetida* Linnaeus, 1758: 582.]

[Note: *mccooki* McCook junior synonym of *foetida* Buckley (synonymy by Wheeler, W.M. 1901b: 520); hence *mccooki* first available replacement name for *foetida* Buckley (Bolton, 1995b: 190).]

Mayr, 1886d: 432 (q.m.).

Combination in *Forelius*: Emery, 1888d: 389.

Junior synonym of *foetida*: Wheeler, W.M. 1901b: 520; Wheeler, W.M. 1902f: 23; Emery, 1913a: 35; Creighton, 1950a: 345; Smith, M.R. 1951a: 836; Kempf, 1972a: 109; Shattuck, 1994: 94.

Status as species: Mayr, 1886d: 432; Cresson, 1887: 257; Emery, 1888c: 362; Dalla Torre, 1893: 168; von Jhering, 1894: 378; Wasmann, 1894: 163; Emery, 1895c: 332; Bruch, 1914: 226; Borgmeier, 1923: 98; Wheeler, W.M. 1942: 215; Bolton, 1995b: 190; Cuezzo, 2000: 249 (redescription); Mackay & Mackay, 2002: 258; Ward, 2005: 9; Guerrero & Fernández, 2008: 58 (in key).

Senior homonym and senior synonym of *maccooki* Forel, 1886b: xxxix.

Senior synonym of *andrei*: Smith, D.R. 1979: 1419; Bolton, 1995b: 190.

maccooki. *Iridomyrmex maccooki* Forel, 1886b: xxxix (w.) U.S.A. (Texas).

[*Iridomyrmex maccooki* Forel, 1878: 382. *Nomen nudum*.]

Status as species: Forel, 1895b: 110; Forel, 1899c: 102; Wheeler, W.M. 1905b: 131; Emery, 1906c: 176; Wheeler, W.M. 1910g: 569; Forel, 1912h: 43; Gallardo, 1916a: 78; Cole, 1937b: 137; Wheeler, W.M. 1938: 252; Shattuck, 1994: 94 (error).

Junior homonym and junior synonym of *mccooki* McCook, 1880: 187.

minor. *Forelius chalybaeus* var. *minor* Forel, 1913l: 241 (w.) ARGENTINA (Mendoza).

Subspecies of *chalybaeus*: Bruch, 1914: 226; Bruch, 1916: 343; Gallardo, 1916a: 86; Santschi, 1916b: 512; Santschi, 1916e: 391; Gallardo, 1919b: 249.

Status as species: Kusnezov, 1957b: 16 (in key); Kempf, 1972a: 109; Shattuck, 1994: 95; Bolton, 1995b: 190.

Junior synonym of *chalybaeus*: Cuezzo, 2000: 239.

modesta. *Forelius nigriventris* var. *modesta* Santschi, 1922b: 374 (w.) ARGENTINA (Formosa).

Subspecies of *nigriventris*: Kempf, 1972a: 110; Shattuck, 1994: 95; Bolton, 1995b: 190.

Junior synonym of *nigriventris*: Cuezzo, 2000: 257.

nigriventris. *Forelius nigriventris* Forel, 1912h: 44 (w.) ARGENTINA (Catamarca).

Brèthes, 1914b: 233 (q.); Cuezzo, 2000: 259 (m.).

Status as species: Forel, 1913l: 241; Forel, 1914d: 287; Brèthes, 1914b: 233; Bruch, 1914: 226; Bruch, 1915: 534; Gallardo, 1915: 22; Gallardo, 1916a: 81; Bruch, 1916: 346; Santschi, 1922b: 374; Kusnezov, 1953b: 339; Kusnezov, 1957b: 17 (in key); Kempf, 1972a: 109; Shattuck, 1994: 95; Bolton, 1995b: 190; Cuezzo, 2000: 257 (redescription); Wild, 2007b: 24.

Senior synonym of *modesta*: Cuezzo, 2000: 257.

Senior synonym of *transiens*: Cuezzo, 2000: 257.

obscurata. *Forelius breviscapus* var. *obscurata* Santschi, 1922b: 374.

[First available use of *Forelius maccooki* r. *breviscapus* var. *obscurata* Forel, 1914d: 286 (w.q.m.) ARGENTINA (Córdoba, Tucumán); unavailable (infrasubspecific) name.]

As unavailable (infrasubspecific) name: Bruch, 1915: 534; Gallardo, 1916a: 81.

Subspecies of *breviscapus*: Kempf, 1972a: 109; Shattuck, 1994: 92; Bolton, 1995b: 190.

Junior synonym of *breviscapus*: Cuezzo, 2000: 235.

paucistricta. *Forelius chalybaeus* var. *paucistricta* Santschi, 1916e: 391 (w.) ARGENTINA (Chubut).

[Misspelled as *paucistriatus* by Kempf, 1972a: 109.]

Subspecies of *chalybaeus*: Kempf, 1972a: 109; Shattuck, 1994: 93; Bolton, 1995b: 190.

Junior synonym of *chalybaeus*: Cuezzo, 2000: 239.

personata. *Forelius chalybaeus* var. *personata* Santschi, 1922b: 375 (w.) ARGENTINA (Córdoba).

Subspecies of *chalybaeus*: Santschi, 1936d: 417; Kempf, 1972a: 109; Shattuck, 1994: 93; Bolton, 1995b: 190.

Junior synonym of *chalybaeus*: Cuezzo, 2000: 239.

pilipes. *Forelius rufus* var. *pilipes* Santschi, 1920d: 382 (w.) ARGENTINA (Santiago del Estero).

[*Forelius rufus* var. *pilipes* Santschi, 1919f: 53. *Nomen nudum*.]

Subspecies of *brasiliensis*: Santschi, 1922b: 374; Kempf, 1972a: 108; Shattuck, 1994: 92; Bolton, 1995b: 190.

Junior synonym of *brasiliensis*: Cuezzo, 2000: 229.

pruinosus. *Tapinoma pruinosum* Roger, 1863a: 165 (w.) CUBA.

Wheeler, G.C. & Wheeler, J. 1951: 185 (l.).

Combination in *Iridomyrmex*: Wheeler, W.M. 1913b: 497;

combination in *Forelius*: Wheeler, G.C. & Wheeler, J. 1986g: 13; Shattuck, 1992c: 95.

Subspecies of analis: Emery, 1913a: 26.

Status as species: Roger, 1863b: 14; Mayr, 1863: 455; Dalla Torre, 1893: 166; Emery, 1895c: 333; Wheeler, W.M. 1905b: 110; Wheeler, W.M. 1905f: 389; Wheeler, W.M. 1910g: 569; Emery, 1913a: 42; Wheeler, W.M. 1913b: 497; Wheeler, W.M. 1913c: 116; Santschi, 1930e: 81; Smith, M.R. 1930a: 5; Wheeler, W.M. 1932a: 12; Wheeler, W.M. 1934g: 178; Cole, 1936a: 37; Cole, 1937b: 136; Cole, 1942: 373; Creighton, 1950a: 342; Smith, M.R. 1951a: 836; Smith, M.R. 1958c: 140; Carter, 1962a: 7 (in list); Smith, M.R. 1967: 364; Kempf, 1972a: 126; Alayo, 1974: 42; Yensen, et al. 1977: 183; Smith, D.R. 1979: 1419; Snelling, R.R. & George, 1979: 166; Petralia & Vinson, 1980: 386; Dlussky, 1981a: 48; Allred, 1982: 478; Wheeler, G.C. & Wheeler, J. 1986g: 57; DuBois & LaBerge, 1988: 144; Deyrup, et al. 1989: 99; Shattuck, 1994: 95; Bolton, 1995b: 190; Cuezzo, 2000: 261 (redescription); Mackay & Mackay, 2002: 258; Deyrup, 2003: 45; Ward, 2005: 9; Coovett, 2005: 108; Guerrero & Fernández, 2008: 57; Branstetter & Sáenz, 2012: 254; Deyrup, 2017: 170; Guerrero, 2019: 710.

Senior synonym of analis: Wheeler, G.C. & Wheeler, J. 1986g: 13; Bolton, 1995b: 190; Coovett, 2005: 108; Ward, 2005: 9.

Senior synonym of testaceus: Snelling, R.R. & George, 1979: 166; Bolton, 1995b: 190; Cuezzo, 2000: 261.

pusillus. *Forelius breviscapus* var. *pusillus* Santschi, 1922b: 374 (w.) ARGENTINA (Córdoba). Cuezzo, 2000: 264 (m.).

Subspecies of breviscapus: Kempf, 1972a: 109; Shattuck, 1994: 93; Bolton, 1995b: 190.

Status as species: Cuezzo, 2000: 263 (redescription); Wild, 2007b: 24; Guerrero & Fernández, 2008: 58 (in key).

Senior synonym of *tucumanus*: Cuezzo, 2000: 263.

***rubriceps*.** *Forelius chalybaeus* var. *rubriceps* Gallardo, 1916a: 85 (date of publication 18.v.1916).

[First available use of *Forelius chalybaeus* r. *grandis* var. *rubriceps* Forel, 1914d: 286 (w.q.m.) ARGENTINA (San Luis); unavailable (infrasubspecific) name.]

[Misspelled as *ruficeps* by Santschi, 1920d: 382.]

As unavailable (infrasubspecific) name: Bruch, 1915: 533.

Subspecies of *chalybaeus*: Bruch, 1916: 345 (date of publication 22.ix.1916); Santschi, 1920d: 382; Santschi, 1929d: 305.

Status as species: Kusnezov, 1957b: 17 (in key); Kempf, 1972a: 110; Shattuck, 1994: 96; Bolton, 1995b: 190; Cuezzo, 2000: 265 (redescription).

ruficeps, misspelling, see under *rubriceps*.

***rufus*.** *Forelius rufus* Gallardo, 1916a: 77 (w.) ARGENTINA (Jujuy).

Santschi, 1922b: 373 (q.m.).

Status as species: Gallardo, 1916c: 261; Santschi, 1920d: 382; Santschi, 1922b: 373; Santschi, 1931e: 281; Kusnezov, 1957b: 17 (in key); Kempf, 1972a: 110; Brandão, 1991: 344; Shattuck, 1994: 96; Bolton, 1995b: 190; Cuezzo, 2000: 268 (redescription).

***symbiotica*.** *Forelius chalybaeus* st. *grandis* var. *symbiotica* Santschi, 1916e: 391 (w.q.) ARGENTINA (Buenos Aires, Rio Negro); unavailable (infrasubspecific) name.

As unavailable (infrasubspecific) name: Gallardo, 1919b: 249.

Declared as unavailable (infrasubspecific) name: Shattuck, 1994: 97.

[*Forelius grandis* var. *symbiotica* Kempf, 1972a: 109. Unavailable name (variety proposed after 1960).]

Material referred to *grandis* by Cuezzo, 2000: 242.

***testaceus*.** *Iridomyrmex pruinosus* var. *testaceus* Cole, 1936b: 121 (w.q.) U.S.A. (Idaho).

Subspecies of *pruinosus*: Cole, 1942: 373; Smith, M.R. 1951a: 836.

Junior synonym of *anale*: Creighton, 1950a: 343; Smith, M.R. 1958c: 140; Smith, D.R. 1979: 1419; Shattuck, 1994: 95.

Junior synonym of *pruinosus*: Snelling, R.R. & George, 1979: 166; Bolton, 1995b: 190; Cuezzo, 2000: 261.

***transiens*.** *Forelius nigriventris* var. *transiens* Santschi, 1922b: 374 (w.) ARGENTINA (Jujuy, Tucumán), BOLIVIA.

Subspecies of *nigriventris*: Kempf, 1972a: 110; Shattuck, 1994: 95; Bolton, 1995b: 190.

Junior synonym of *nigriventris*: Cuezzo, 2000: 257.

***tucumanus*.** *Neoforelius tucumanus* Kusnezov, 1953b: 330, figs. 1-12 (w.q.m.) ARGENTINA (Tucumán).

Combination in *Forelius*: Shattuck, 1992c: 95.

Status as species: Kempf, 1972a: 160; Shattuck, 1994: 96; Bolton, 1995b: 190.

Junior synonym of *pusillus*: Cuezzo, 2000: 263.