

**Chromosome Observations on Tropical Ants from Western  
Malaysia. II**

Hirotami T. IMAI, William L. BROWN Jr.<sup>1)</sup>, Masao KUBOTA<sup>2)</sup>,  
Hoi-Sen YONG<sup>3)</sup> and Yow Pong THO<sup>4)</sup>

Imai and Kubota went to West Malaysia in 1982 and 1983 for a karyological survey of tropical ants. The project was performed in cooperation with H. S. Yong and Y. P. Tho. The identification of species was made by M. Kubota and W. L. Brown, though it remains unsettled yet in some ant groups, e.g., *Pheidole* and *Camponotus*. We already described chromosomes of 73 ant species collected in 1982 (Ann. Rep. Natl. Inst. Genetics, **32**: 71–73). The present paper reports chromosomes of 86 species (39 genera, 6 subfamilies) collected in 1983. A total of 119 colonies, labelled as HI83-(1-119), were collected from the following localities: campus of Forest Research Institute (HI83-1–42), Jeram Toi (HI83-43–46, 49), Pasoh Forest (HI83-47–74, 103), Gombak Station of the University of Malaya (HI83-76–83, 96, 109–115), Genting Highlands (HI83-84–95), Ulu Gombak

<sup>1)</sup> Department of Entomology, Cornell University, Ithaca, New York, 14853, U.S.A.

<sup>2)</sup> Nakasone 13, Odawara, Kanagawa-ken, 250, Japan.

<sup>3)</sup> Department of Genetics and Cellular Biology, University of Malaya, Kuala Lumpur, Malayasia.

<sup>4)</sup> Forest Research Institute, Kepong, Malaysia.

Table 1. Chromosome numbers of Malaysian ants. II

Taxa (Colony number)	Chrom. number (n) 2n	Taxa (Colony number)	Chrom. number (n) 2n
PONERINAE		DORYLINAЕ	
<i>Platythyrea quadridenta</i> (HI83-22, 50)	(9) 18	<i>Aenictus laeviceps?</i> (HI83-103)	22
<i>P. tricuspidata</i> (HI83-116)	96	MYRMICINAE	
<i>Gnamptogenys menadensis*</i> (HI83-61)	42	<i>Aphaenogaster beccarii*</i> (HI83-55)	30
<i>G. binghami</i> (HI83-21)	(22)	<i>Pheidole sp. I*</i> (HI83-101)	(10)
<i>Leptogenys diminuta=sp. I*</i> (HI83-14, 15, 86)	38	<i>P. sp. 7</i> (HI83-7)	16
<i>L. myops=sp. 2*</i> (HI83-113)	(24)	<i>P. sp. 11</i> (HI83-53)	22
<i>L. sp. 3</i> (HI83-109)	48	<i>P. sp. 12</i> (HI83-44)	(10) 20
<i>L. iridescent</i> (HI83-47, 65)	46	<i>P. sp. 13</i> (HI83-102)	38
<i>L. borneensis</i> (HI83-66)	46	<i>P. sp. 14</i> (HI83-32)	20
<i>Diacamma sp. I*</i> (HI83-16, 99, 106)	44	<i>P. sp. 15</i> (HI83-6)	20
<i>Amblyopone reclinata</i> (HI83-20)	38	<i>P. sp. 16</i> (HI83-104)	(9) 18
<i>Pachycondyla "n. sp. A of Brown"</i> = <i>Mesoponera sp. I</i> (HI83-63, 71)	(18) 36	<i>P. sp. 17</i> (HI83-27)	20, 21
<i>P. leeuwenhoekii</i> (8) 16		Translocation polymorphism	
<i>P. rubra*=Mesoponera sp. 2</i> (HI83-3, 11, 12)	38, 40	<i>P. sp. 18</i> (HI83-8, 88)	20
Robertsonian polymorphism		<i>P. sp. 19</i> (HI83-9, 90)	20
<i>P. tridentata</i> (HI83-51)	28	<i>P. sp. 20</i> (HI83-58, 60, 108)	(18) 36
<i>Ponera japonica</i> (HI83-94)	12	<i>P. megacephala</i> (HI83-82)	20
<i>Hypoponera pruinosa</i> (HI83-110)	(12) 24	<i>Crematogaster sp. I*</i> (HI83-59)	26
<i>H. sp. 2</i> (HI83-35)	38	<i>C. sp. 4*</i> (HI83-5, 49)	24
<i>H. sp. 3</i> (HI83-45)	36	<i>C. sp. 5</i> (HI83-4)	26
<i>Cryptopone testacea</i> (HI83-54)	(9) 18	<i>C. sp. 6</i> (HI83-72)	26
<i>Odontomachus simillimus</i> = <i>sp. I*</i> (HI83-17, 18)	44	<i>Monomorium sp. 2*</i> (HI83-85)	22
<i>O. rixosus=sp. 2*</i> (HI83-38)	(15) 30	<i>M. sp. 3</i> (HI83-96)	(11) 22
<i>O. latidens</i> (HI83-25)	(15)	<i>M. sp. 4</i> (HI83-105)	(11)
PSEUDOMYRMECINAE		<i>Lophomyrmex bedoti</i> (HI83-73, 75)	38
<i>Tetraponera sp. 2</i> (HI83-97)	42	<i>Vollenhovia sp. I</i> (HI83-46, 69)	49, 50
		Robertsonian polymorphism	
		<i>V. sp. 2</i> (HI83-93, 95)	33, 34
		Robertsonian polymorphism	
		<i>Acanthomyrmex sp. 3</i> (HI83-10, 98, 114)	(11) 22
		<i>Pristomyrmex sp. 2</i> (HI83-77)	(14)

Table 1. Continued

Taxa (Colony number)	Chrom. number (n) 2n	Taxa (Colony number)	Chrom. number (ff) 2n
<i>Lordomyrma</i> sp. I (HI83-37)	(11) 22	<i>Plagiolepis</i> sp. I (HI83-62)	18
<i>Myrmicaria</i> sp. 2* (HI83-87)	44	<i>Pseudolasius</i> sp. I (HI83-57, 70)	( 8)
<i>M.</i> sp. 4 (HI83-43)	(23) 46	<i>P.</i> sp. 2 (HI83-67, 68)	(15, 17,
<i>Tetramorium pnyxis</i> (HI83-56)	20	B-chromosome	19) 30
<i>T. eleates</i> (HI83-30)	28	polymorphism	
<i>T. seneb</i> ? (HI83-29)	20	<i>Camponotus festinus</i> * (HI83-24, 52)	(19) 38
<i>Smithistruma taipingensis</i> * (HI83-111)	(12) 24	<i>C.</i> sp. 2* (HI83-19)	40
<i>Strumigenys godeffroyi</i> (HI83-31)	40	<i>C.</i> sp. 7 (HI83-107)	(20)
<i>Dacetinops concinna</i> (HI83-115)	16	<i>C.</i> sp. 8 (HI83-23)	38, 39
<i>Eurhopalothrix</i> n. sp. ( <i>procera</i> group)	18	Robertsonian polymorphism	
Brown (in press)		<i>C.</i> sp. 9 (HI83-80, 81)	40
DOLICHODERINAE		<i>C.</i> sp. 10 (HI83-41, 42)	(13, 14)
<i>Dolichoderus bituberculatus</i> * (HI83-83) B-chromosome	30, 31, 32, 33	Robertsonian polymorphism	
polymorphism		<i>C.</i> sp. 11 (HI83-84)	52
<i>Iridomyrmex cordatus</i> (HI83-74)	16	<i>C.</i> sp. 12 (HI83-100)	18
<i>Technomyrmex</i> sp. 1* (HI83-118)	30	<i>C.</i> sp. 13 (HI83-78)	18
FORMICINAE		<i>Polyrhachis illaudata</i> (HI83-36)	(18)
<i>Anopolepis longipes</i> * (HI83-1, 2, 117)	34	<i>P. hector</i> (HI83-76, 112)	(21) 42
<i>Acropyga acutiventris</i> * (HI83-28, 119)	28, 29	<i>P. rastellata</i> (HI83-39, 40, 79)	42
Robertsonian polymorphism		<i>Echinopla</i> sp. 1 (HI83-48, 64)	(12) 24
		<i>Paratrechina</i> sp. 4* (HI83-34)	16
		<i>P.</i> sp. 6 (HI83-92)	30
		<i>P.</i> sp. 7 (HI83-33)	16
		<i>Prenolepis jerdoni</i> * (HI83-26, 91)	(16, 20, 25, 27)
		B-chromosome	30, 31,
		polymorphism	32, 34

N.B. The species with asterisks were observed also in the survey of 1982.

(HI83-97-108), and Templer Park (HI83-116-119). Among 119 colonies, we were able to observe chromosome spreads in all samples except HI83-13, 46, 57, 63, 79, 89, and 119. The results are summarized in Table 1. In addition, two species collected at Sungey Menala Reserve, Negeri Sembilan, in 1981 by Brown and Pong (*Pachycondyla leeuwenhoeki* and *Eurhopalothrix*

*n. sp.*), were karyotyped separately by Brown; these have been added to Table 1.