

## **TERATANER**

- alluaudi.** *Atopomyrmex alluaudi* Emery, 1895f: 341, fig. 2 (w.) MADAGASCAR.  
Wheeler, G.C. & Wheeler, J. 1980: 536 (l.); Alpert, 1992: 123 (ergatoid q., m.).  
Combination in *Terataner*: Emery, 1912b: 103.  
**Status as species:** Wasmann, 1897: 250; Emery, 1899f: 274; Emery, 1912b: 103; Wheeler, W.M. 1922a: 1028; Emery, 1924d: 242; Bolton, 1981b: 296; Alpert, 1992: 119; Bolton, 1995b: 403; Hita Garcia, Fischer, *et al.* 2017a: 11 (in key).
- balrog.** *Terataner balrog* Hita Garcia, in Hita Garcia, Fischer, *et al.* 2017a: 12, figs. 1, 3B, 3D, 4-7, S1 and S2 (w.ergatoid q.) MADAGASCAR.
- bottegoi.** *Atopomyrmex bottegoi* Emery, 1896i: 155, fig. (w.) SOMALIA.  
Type-material: holotype worker.  
Type-locality: Somalia: Lugh (V. *Bottego*).  
Type-depository: MSNG.  
Combination in *Terataner*: Emery, 1912b: 103;  
combination in *Terataner (Tranetera)*: Arnold, 1952b: 130.  
**Status as species:** Emery, 1912b: 103; Santschi, 1914b: 72; Wheeler, W.M. 1922a: 885; Emery, 1924d: 242; Menozzi, 1930b: 109; Bolton, 1981b: 291 (redescription); Bolton, 1995b: 403; Hita Garcia, *et al.* 2013: 217.  
Distribution: Kenya, Somalia, Tanzania.
- elegans.** *Terataner elegans* Bernard, 1953b: 243, fig. 13A (q.) GUINEA.  
Type-material: holotype queen.  
Type-locality: Guinea: Kéoulenta (*Lamotte*).  
Type-depository: MNHN.  
Bolton, 1981b: 292 (w.).  
**Status as species:** Bolton, 1981b: 292 (redescription); Bolton, 1995b: 403.  
Distribution: Ghana, Guinea, Nigeria.
- foreli.** *Atopomyrmex foreli* Emery, 1899f: 274, figs. (w.m.) MADAGASCAR.  
Emery, 1899f: 274 (m.).  
Combination in *Terataner*: Emery, 1912b: 103.  
**Status as species:** Emery, 1899f: 274; Emery, 1912b: 103; Wheeler, W.M. 1922a: 1029; Emery, 1924d: 242; Bolton, 1981b: 296; Alpert, 1992: 119; Bolton, 1995b: 403; Hita Garcia, Fischer, *et al.* 2017a: 11 (in key).
- luteus.** *Atopomyrmex luteus* Emery, 1899e: 477 (w.) CAMEROON.  
Type-material: holotype worker.  
Type-locality: Cameroon: (no further data) (*Conradi*).  
Type-depository: MSNG.  
[Note: specimen labelled “type” in MCZC is not mentioned in the original description.]  
Combination in *Terataner*: Emery, 1912b: 103.  
**Status as species:** Stitz, 1910: 145; Emery, 1912b: 103; Wheeler, W.M. 1922a: 885; Emery, 1924d: 242; Bolton, 1981b: 293 (redescription); Bolton, 1995b: 403.  
Distribution: Cameroon, Ghana.
- nymeria.** *Terataner nymeria* Hita Garcia, in Hita Garcia, Fischer, *et al.* 2017a: 21, figs. 1, 3C, 3F-G, 8-10, S3 (w.) MADAGASCAR.
- piceus.** *Terataner piceus* Menozzi, 1942: 173 (w.) EQUATORIAL GUINEA.  
[Also given as new in Eidmann, 1944: 469 (in list).]  
Type-material: holotype worker.  
Type-locality: Equatorial Guinea (“Guinea Spagnola”): Rio Benito, 1939-40 (*H. Eidmann*).  
Type-depository: holotype not found in IEUB, presumed lost (Bolton, 1981b: 293).  
**Status as species:** Bolton, 1981b: 293 (redescription); Bolton, 1995b: 403.  
Distribution: Cameroon, Equatorial Guinea, Ghana, Nigeria.
- rufipes.** *Terataner rufipes* Emery, 1912b: 104 (w.) MADAGASCAR.  
**Status as species:** Wheeler, W.M. 1922a: 1029; Emery, 1924d: 242; Bolton, 1981b: 296; Bolton, 1995b: 403; Hita Garcia, Fischer, *et al.* 2017a: 11 (in key).

- scotti.** *Atopomyrmex scotti* Forel, 1912k: 160 (w.m.) SEYCHELLES IS (Praslin I., Silhouette I.).  
[*Terataner scoti* Emery, 1912b: 103. *Nomen nudum* (attributed to Forel).]  
Combination in *Terataner*: Wheeler, W.M. 1922a: 1029.  
**Status as species:** Emery, 1912b: 103; Wheeler, W.M. 1922a: 1029; Emery, 1924d: 242; Bolton, 1981b: 296; Bolton, 1995b: 403; Dorow, 1996a: 81.
- steinheili.** *Atopomyrmex steinheili* Forel, 1895h: 485 (w.) MADAGASCAR.  
Combination in *Terataner*: Emery, 1912b: 103.  
**Status as species:** Emery, 1912b: 103; Wheeler, W.M. 1922a: 1029; Emery, 1924d: 242; Bolton, 1981b: 296; Bolton, 1995b: 403; Hita Garcia, Fischer, *et al.* 2017a: 11 (in key).
- transvaalensis.** *Terataner (Tranetera) transvaalensis* Arnold, 1952b: 130, 2 figs. (w.) SOUTH AFRICA.  
Type-material: holotype worker, 1 paratype worker.  
Type-locality: South Africa: eastern Transvaal, Marieps (or Mariepskop), v.1951 (*J.C. Faure*).  
Type-depository: SAMC (holotype).  
**Status as species:** Bolton, 1981b: 294 (redescription); Bolton, 1995b: 403.  
Distribution: South Africa.
- velatus.** *Terataner velatus* Bolton, 1981b: 295 (w.) GHANA, IVORY COAST.  
Type-material: holotype worker, 5 paratype workers.  
Type-locality: Ghana: Okumaning, 12.viii.1969, ant ecology sample K12 (*D. Leston*), 1 paratype same data but sample K11; non-type-locality paratypes: 1 paratype Ghana: Kade, ant ecology sample K96 (*D. Leston*), 2 paratypes Ghana: Enchi, 17.v.1969, ant ecology sample E5-8 (*D. Leston*), 1 paratype Ivory Coast: Banco Forest near Abidjan, 10.i.1963, no. A30 (*W.L. Brown*).  
Type-depositories BMNH (holotype); BMNH, MCZC (paratypes).  
**Status as species:** Bolton, 1995b: 403.  
Distribution: Ghana, Ivory Coast.
- xaltus.** *Terataner xaltus* Bolton, 1981b: 296, fig. 52 (w.) MADAGASCAR.  
Type-material: holotype worker.  
Type-locality: Madagascar: Forêt de Zombitsy, near Sakaraha, 650 m, 16.xii.1959 (no collector's name).  
Type-depository: MCZC.  
**Status as species:** Bolton, 1995b: 403; Hita Garcia, Fischer, *et al.* 2017a: 11 (in key).  
Distribution: Madagascar.